

Original Article

Depressive symptoms and psychosocial risk factors in high complexity obstetric patients admitted to a critical care obstetric unit in Cali, Colombia[☆]



Ana María Guerra, Diana María Dávalos Pérez, Alejandro Castillo Martínez*

Fundación Clínica Valle del Lili, Universidad ICESI, Cali, Colombia

ARTICLE INFO

Article history:

Received 10 August 2015

Accepted 5 September 2016

Available online 18 October 2017

Keywords:

Depresión

Perinatal

Screening

ABSTRACT

Depression is the main cause of years lost due to disability in the world, and it affects 50% more women 50% than men. Perinatal depression has been linked with more anxiety, a chronic course, and disability than depression in other life stages. In spite of its high prevalence and serious health effects on both mother and foetus, it is frequently underdiagnosed. This study was performed on all high risk obstetric patients admitted to a critical care obstetric unit in Cali, Colombia, from January to June, 2014. Depressive symptoms and psychosocial risk factors were screened by means of a survey and the Edinburgh Prenatal Depression Scale (EPDS). A total of 695 women were included, of whom 30.2% had depressive symptoms on the EPDS, and 3.6% reported having self-injury thoughts in the last 7 days. Our findings are consistent with previous reports on a history of child abuse and family depression as risk factors. It was also suggested that lower socio-economic status is associated with more vulnerability. Early screening is needed to ensure timely detection and treatment.

© 2016 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Detección de síntomas depresivos en mujeres gestantes de alta complejidad obstétrica y factores correlacionados

RESUMEN

La depresión es la principal causa de años perdidos por discapacidad en el mundo, y en mujeres la carga de la enfermedad es un 50% mayor que en varones. La depresión perinatal tiene mayor asociación con ansiedad, cronicidad y discapacidad que en otras etapas de la vida y es poco reconocida a pesar de su alta prevalencia y sus graves efectos en la salud de

Palabras clave:

Depresión

Perinatal

Detección

DOI of original article: <http://dx.doi.org/10.1016/j.rcp.2016.09.002>.

[☆] Please cite this article as: Guerra AM, Dávalos Pérez DM, Castillo Martínez A. Detección de síntomas depresivos en mujeres gestantes de alta complejidad obstétrica y factores correlacionados. Rev Colomb Psiquiat. 2017;46:215-221.

* Corresponding author.

E-mail address: acastillo@fcvl.org (A. Castillo Martínez).

<http://dx.doi.org/10.1016/j.rcpeng.2017.09.009>

2530-3120/© 2016 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

la madre y el feto. Se realizó un estudio en pacientes ingresadas y valoradas en una unidad de alto riesgo obstétrico en el proceso de implementación del programa durante los meses de enero a junio de 2014; se utilizaron la *Edinburgh Prenatal Depression Scale* (EPDS) y un cuestionario previamente desarrollado para identificar la presencia de síntomas depresivos y factores de riesgo psicosocial. Se evaluó sistemáticamente a más de 600 mujeres, y se encontró que el 30,2% de las mujeres evaluadas tenían síntomas depresivos según la EPDS y el 3,6% había tenido ideas de autolesión en los últimos 7 días. Los factores relacionados evaluados indican una condición de vulnerabilidad de las mujeres con menor nivel socioeconómico. Los datos encontrados reiteran la importancia de detectar factores que tradicionalmente el personal de salud no identifica ni interviene a lo largo de la vida y particularmente durante la gestación.

© 2016 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Depression is the most prevalent mental illness and the leading cause of disability-adjusted life years worldwide.¹ Depression is twice as prevalent among women than men. Prevalence peaks during reproductive age, the clinical course varies greatly, and there is a high risk of recurrence.

Perinatal depression is defined as an episode of major depressive disorder occurring either during pregnancy or within the first 12 months postpartum; it is more closely associated with anxiety, chronicity and disability than depression in other stages of life.²⁻⁴

Prenatal depression in particular, despite its high prevalence and serious effects on the health of the mother and the foetus, is a poorly recognised clinical entity. It increases the risk of both adverse perinatal outcomes and alterations in the development and health of the foetus until adulthood.⁵⁻¹³ Likewise, depression during pregnancy is the main predictor of postpartum depression and life-long recurrence of depressive episodes in women.^{14,15}

Prenatal depression is a risk factor for miscarriage, preeclampsia, preterm birth and low birth weight. It is also a risk factor for the child's health, and is associated with delayed psychomotor development, child abuse, depression, attention and behavioural disorders, dissociative disorder and poor academic performance.¹⁶⁻²⁶

In Latin America, the prevalence of depression during pregnancy is estimated to be between 17.4% and 40.1%.²⁷⁻²⁹ In Colombia, estimates suggest that 32.8% of pregnant adolescents suffer from depression.³⁰

Studies have identified a series of risk factors for perinatal depression: personal and family history of mental illness, stressful life events in the year before pregnancy, conjugal conflict, domestic violence, single motherhood, social and economic deprivation, low educational level, unemployment, history of abuse during childhood, inadequate social support, young maternal age and loss of or separation from a parent before 16 years of age.^{27,30-39} Some studies have also observed a certain interaction between these risk factors, suggesting that the higher the number of risk factors, the greater the severity of depressive symptoms.⁴⁰

The protective factors so far identified are: higher income, higher educational level, permanent or secure job, partner with a job, and adequate emotional and social support.^{28,35}

Different scales can be used to detect symptoms of perinatal depression. The Edinburgh Postnatal Depression Scale (EPDS), for example, is a short, self-administered 5-10 min scale designed to improve detection of postpartum depression in rural communities in developing countries, and has also been tested for use in the prenatal period.

The first step in treating perinatal depression and preventing negative outcomes in both the mother and baby is to introduce mechanisms for the early detection of depression or risk factors for depression in pregnant women.

Methods

We performed a cross-sectional study in patients admitted to the High Complexity Obstetric Care (HCOC) unit from January to June 2014, during the introduction of the programme. We used the EPDS and an in-house questionnaire to identify the presence of depressive symptoms and psychosocial risk factors.

Our aim was to detect the presence of depressive symptoms using the EPDS and the frequency of the psychosocial variables routinely evaluated by the interdisciplinary team in patients with depressive symptoms detected by the EPDS.

The EPDS scale consists of 10 items; the first 9 explore the cognitive symptoms of depression and the last item is specific for self-harm ideas in the last 7 days. It has been translated into Spanish and validated in Latin America in general, and Colombia in particular.⁴¹ With a cut-off point of 10, the EPDS has a sensitivity of 100%, a specificity of 80% and a positive predictive value of 37%.⁴² The EPDS is a screening scale, and it was initially administered by the study nurse at admission, together with a questionnaire developed by the authors to evaluate certain psychosocial and clinical variables.

In this study, we present the results of our evaluation of the data from the first stage of the implementation of a programme for early detection and comprehensive management of depressive symptoms and depression in pregnant women treated in the HCOC unit of the Fundación Valle del Lili. The unit is staffed by a multidisciplinary team that

includes gynaecologists, psychiatrists, psychologists, occupational therapists and social workers. The sample comprised all patients aged over 18 years admitted to the HCOG hospitalisation unit for gynaecological disorders during pregnancy or after delivery during the study period who agreed to participate in the study and signed the informed consent form.

The study was conducted according to international bioethical standards, and was approved by the ethics committee of the Fundación Valle de Lili Clinical.

Although this is an administrative database, this study was planned from the moment data collection started. The Fundación Valle del Lili is a private tertiary referral clinic located in the south west of Colombia. It provides highly complex medical care for individuals included in both the contributory and subsidised schemes of the General Social Security System (GSSS).

Analytical plan

We conducted an initial exploratory analysis and a descriptive analysis of the population, and determined the frequency of depressive symptoms in the sample of women and the distribution of sociodemographic variables. We then compared the distribution of study variables between groups using the χ^2 and Fisher's exact test, as appropriate. For normally distributed continuous variables the Student's t-test was used; for variables with non-normal distribution, the Mann-Whitney U test was used. The strength of association between the dependent variable and independent variables (bivariate analysis) was determined by the odds ratio (OR) and 95% confidence interval (95% CI).

Results

A total of 695 women admitted to the HCOG during the period from January to June 2014 were evaluated. The average age was 27.4 years; 40% were housewives and 67% were enrolled in the contributory scheme of the GSSS (Tables 1 and 2).

The mean age of women enrolled in the subsidised scheme was lower ($p=0.0001$) than that of those enrolled in the contributory regime (Tables 3 and 4).

Of the total sample, 77% were pregnant at admission, 54% of the women evaluated had 1 or 2 children, and 96% of them had received prenatal care. It was found that 30.2% of the women scored ≥ 10 on the EPDS, and 3.6% reported thoughts of self-harm (Tables 1–4).

The psychosocial risk factors evaluated were age, health insurance (a proxy for social and economic status, since individuals enrolled in the contributory regimen have a higher level of income than those in the subsidised regimen), number of children, history of abortion, history of sexual and emotional abuse, previous suicide attempts, personal or family history of depression and cohabitation with the spouse (Table 1).

The bivariate analysis showed that women with depressive symptoms were more likely to report sexual (OR=6.6; 95% CI, 1.93–28.76) or emotional (OR=3.4, 95% CI, 1.37–8.64) abuse at

Table 1 – Sociodemographic characteristics.

Patients (n)	695
<i>Age groups</i>	
12–18 years	80 (11.5)
19–29 years	342 (49.2)
30–39 years	238 (34.3)
≥ 40 years	35 (5.0)
<i>Health insurance</i>	693 (100)
Subsidised	179 (25.8)
Contributory	464 (67.0)
Prepaid healthcare and other regimens	50 (7.2)
<i>Occupation/education</i>	695 (100)
Housewife	281 (40.3)
Professional	108 (15.6)
Technician	85 (12.2)
Student	68 (9.8)
Other	68 (9.8)
Employed	45 (6.5)
No data	40 (5.8)
<i>Condition at admission</i>	686 (100)
Pregnancy	532 (77.5)
Postpartum	154 (22.5)
<i>Gestation, n</i>	357
First trimester	21
Second trimester	69
Third trimester	267
<i>Number of children</i>	659 (100)
None	238 (36.1)
1–2	357 (54.2)
3–4	53 (8.0)
5–12	11 (1.7)
<i>History of abortion</i>	695 (100)
None	504 (72.5)
1–2	171 (24.6)
3–5	15 (2.2)
No data	5 (0.7)
<i>Other history</i>	348
Caesarean	284 (40.9)
Stillbirth	46 (6.6)
Ectopic	18 (2.6)

The values are expressed as n (%) unless otherwise indicated.

some time during their life, previous suicide attempts and a personal and/or family history of depression (Table 5).

Discussion

In this study, we present the data collected following the introduction of a programme for screening and treatment of depression in women with highly complex pregnancies at the Fundación Valle del Lili. It is the first study of its kind to be conducted in this population in Colombia. We found that 30.2% of study subjects presented depressive symptoms on the EPDS, and 3.6% reported thoughts of self-harm in the last 7 days.

In their systematic review, Gavin et al. showed that the combined prevalence of major and minor depression reported

Table 2 – Clinical variables.

Edinburgh Postnatal Depression Scale > 10	695 (100)
Yes	210 (30.2)
No	485 (60.8)
Follow-up	695 (100)
Yes	232 (33.4)
No	463 (66.6)
Prenatal care	692 (100)
Yes	665 (96.1)
No	27 (3.9)
Treatment by interdisciplinary team	63
Psychiatrist	47
Psychologist	8
Social worker	3
Occupational therapist	5
Report of suicidal ideation	695 (100)
Yes	25 (3.6)
No	670 (96.4)

Table 3 – Age according to health insurance.

	Patients, n	Age (years)	
		Mean \pm SD	Median (range)
Total	695	27.4 \pm 7.2	27 (12–52)
Health Insurance			
Subsidised	179	25 ^a \pm 7.9	23 (12–51)
Contributory	464	28.1 ^a \pm 6.8	28 (14–38)
Other	50	29.6 ^a \pm 5.4	30 (19–38)

^a Kruskal–Wallis = 32.464 with 2 d.f. $p = 0.0001$.

Table 4 – Depressive symptoms according to the Edinburgh scale.

	EPDS > 10	EPDS < 10
Age groups, n	210	485
12–18 years	31 (38.8)	49 (61.2)
19–29 years	100 (29.2)	242 (70.8)
30–39 years	65 (27.3)	173 (72.7)
≥ 40 years	14 (40)	21 (60)
Health insurance, n	210	483
Subsidised	59 (33)	120 (67)
Contributory	138 (29.7)	326 (70.3)
Prepaid healthcare and other regimens	13 (26)	37 (74)

EPDS: Edinburgh Postnatal Depression Scale.

in the studies included in their meta-analysis varied from 6.5% to 12.9% in the different trimesters of pregnancy. However, the published rate of prenatal depression in Latin America is higher. Studies in Mexico and Chile have shown a prevalence of prenatal depression confirmed by clinical interview of 17.4% and 22%, respectively.²⁷

In Colombia, studies using the Zung scale reported a prevalence of depressive symptoms in pregnant adolescents of 32.8%,³⁰ while in Ecuador, the CESD-10 (Center for Epidemiologic Studies Depression) scale showed that 56.6% of pregnant women presented depression.⁴³

Other researchers have used the EPDS with higher cut-off points to diagnose depression. In Peru, Lam et al. reported that 34.1% of pregnant women had an EPDS score ≥ 14 , and Luna-Mattos et al. found 40.1% with an EPDS score ≥ 13 . In Colombia, Borda-Pérez et al. reported a prevalence of 19.9% with an EPDS score ≥ 13 and Peinado-Valencia et al. found 31.5% with an EPDS score ≥ 14 .

This shows that the prevalence of probable depression found in our study population is in line with national and South American data, and is greater than that described in other parts of the world.

A review of the studies included in the databases analysed failed to identify any papers dealing with the systematic evaluation and treatment of depressive symptoms in this patient group in Colombia.

In our study, the GSSS regimen of each patient was used as a proxy for social and economic status, as enrolment in this regimen means that the woman or her partner are currently employed, while individuals that are unemployed and have no private means receive medical attention under the subsidised regimen. For example, the differences found in age and number of children show that women from a lower social and economic status are more at risk of depression in this population.

The finding that 12% of all women with a positive EPDS score had ideas of self-harm is in line with the expected rate of suicidal ideation in the depressed patient population, which could be explained by the severity of the depressive symptoms found in this population.

Our findings highlight the importance of detecting risk factors that healthcare personnel do not usually identify and, therefore, do not treat, particularly during pregnancy. As previously mentioned, previous episodes of depression and a history of abuse and sexual violence increase the risk for the health and well-being of women and their children.

Some studies have used other variables not included in our study to evaluate this problem. For example, Peinado Valencia et al. found the number of children and non-Catholic religion to be risk factors; Borda et al. found an association with a history of abortion (OR = 3.39, 95% CI, 1.41–8.15), some degree of family dysfunction (OR = 2.78, 95% CI, 1.08–7.1) and activity outside the home (OR = 2.92, 95% CI, 1.03–8.23).

In Mexico, Lara et al. found a history of depression in the mother of the pregnant woman (risk increased 0.8-fold), personal history of depression (risk increased 1.08-fold), lack of social support (risk increased 1.7-fold) and no partner (risk increased 1.51-fold) to be significant prenatal depression variables. These findings are consistent with our study.

Although probabilistic sampling was not performed, more than 600 women were systematically evaluated in this study. This provided a larger sample size than previous studies with similar methodologies conducted at a national and regional level. In addition, being a tertiary referral hospital, our sample is made up of women referred to the Valle de Lili Clinic from different public and private hospitals throughout the south-

Table 5 – Bivariate analysis.

	EPDS > 10 (n = 210)	EPDS < 10 (n = 485)	OR	95% CI	p
<i>Age groups</i>					
12–18 years	31 (38.8)	49 (61.2)	1.53	0.89–2.61	0.09
19–29 years*	100 (29.2)	242 (70.8)			
30–39 years	65 (27.3)	173 (72.7)	0.9	0.62–1.33	0.61
≥40 years	14 (40)	21 (60)	1.61	0.73–3.48	0.19
<i>Health insurance</i>					
Subsidised	59 (33)	120 (67)	1.16	0.79–1.71	0.42
Contributory*	138 (29.7)	326 (70.3)			
Prepaid healthcare and other regimens	13 (26)	37 (74)	0.83	0.39–1.66	0.58
<i>History of sexual abuse</i>					
Yes	11	4	6.62	1.93–28.76	<0.01
No	199	479			
<i>History of emotional abuse</i>					
Yes	14	196	3.38	1.37–8.64	<0.01
No	10	473			
<i>History of physical abuse</i>					
Yes	5	17	0.67	0.19–1.91	0.43
No	205	466			
<i>Living with spouse</i>					
Yes	102	204	1.29	0.92–1.81	0.12
No	108	279			
<i>History of depression</i>					
Yes	33	16	5.44	2.82–10.83	<0.01
No	177	467			
<i>Family history of depression</i>					
Yes	23	16	3.59	1.77–7.43	<0.01
No	187	467			
<i>History of attempted suicide</i>					
Yes	10	1	24.1	3.03–189.46	<0.01
No	200	482			

EPDS: Edinburgh Postnatal Depression Scale.

west region of Colombia. The methodology used to select study subjects, however, exposes the study to memory bias, insofar as patients with depressive symptoms are more likely to remember traumatic events in their lifetime than other, non-depressive patients.

This study provides some data that may be useful in future prospective studies that use the clinical interview to detect depressive disorders, and which evaluate specific outcomes – miscarriage, preeclampsia, preterm birth and low birth weight – associated with prenatal depression.^{15–18}

This study also contributes to public health, insofar as it shows the need to improve the screening of depressive symptoms and the treatment of depression in high-risk pregnant women in Colombia, and can improve the health prognosis of both these patients and their children.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. Surkan PJ, Kennedy CE, Hurley KM, Black MM. Maternal depression and early childhood growth in developing countries: systematic review and meta-analysis. *Bull World Health Organ.* 2011;89:608–15.
2. Meltzer-Brody S. New insights into perinatal depression: pathogenesis and treatment during pregnancy and postpartum. *Dial Clin Neurosci.* 2011;13:89–100.

3. Cooper P, Murray L. Course and recurrence of postnatal depression. Evidence for the specificity of the diagnostic concept. *Br J Psychiatry*. 1995;166:191-5.
4. Patel V, Rodrigues M, DeSouza N. Gender, poverty, and postnatal depression: a study of mothers in Goa, India. *Am J Psychiatry*. 2002;159:43-7.
5. Monk C, Spicer J, Champagne F. Linking prenatal maternal adversity to developmental outcomes in infants: the role of epigenetic pathways. *Dev Psychopathol*. 2012;24:1361-76.
6. Van Den Bergh B, Mennes M, Oosterlaan J, Stevens V, Stiers P, Marcoen A, et al. High antenatal maternal anxiety is related to impulsivity during performance on cognitive tasks in 14-15 year olds. *Neurosci Biobehav Rev*. 2005;29:259-69.
7. Evans J, Melotti R, Heron J, Ramchandani P, Wiles N, Murray L, et al. The timing of maternal depressive symptoms and child cognitive development: a longitudinal study. *J Child Psychol Psychiatry*. 2011;53:632-40.
8. Koutra K, Chatzi L, Bagkeris M, Vassilaki M, Bitsios P, Kogevinas M. Antenatal and postnatal maternal mental health as determinants of infant neurodevelopment at 18 months of age in a mother-child cohort (Rhea Study) in Crete, Greece. *Soc Psychiatry Psychiatr Epidemiol*. 2012;48:1335-45.
9. Hay D, Pawlby S, Waters C, Sharp D. Antepartum and postpartum exposure to maternal depression: different effects on different adolescent outcomes. *J Child Psychol Psychiatry*. 2008;49:1079-88.
10. Leitner Y, Fattal-Valevski A, Geva R, Eshel R, Toledano-Alhadeef H, Rotstein M, et al. Neurodevelopmental outcome of children with intrauterine growth retardation: a longitudinal, 10-year prospective study. *J Child Neurol*. 2007;22:580-7.
11. Alexander GR. Prematurity at birth: determinants, consequences and geographic variation. Preterm birth. Causes, consequences and prevention. Washington: IOM of the National Academy Press; 2006.
12. Field T, Diego M, Hernandez-Reif M. Prenatal depression effects on the fetus and newborn: a review. *Infant Behav Dev*. 2006;29:445-55.
13. Alder J, Fink N, Bitzer J, Hösl I, Holzgreve W. Depression and anxiety during pregnancy: a risk factor for obstetric, fetal and neonatal outcome? A critical review of the literature. *J Matern Fetal Neonatal Med*. 2007;20:189-209.
14. Rahman A, Creed F. Outcome of prenatal depression and risk factors associated with persistence in the first postnatal year: prospective study from Rawalpindi, Pakistan. *J Affect Disord*. 2007;100:115-21.
15. Pawlby S, Hay D, Sharp D, Waters C, O'Keane V. Antenatal depression predicts depression in adolescent offspring: prospective longitudinal community-based study. *J Affect Disord*. 2009;113:236-43.
16. Nakano Y, Oshima M, Sugiura-Ogasawara M, Aoki K, Kitamura T, Furukawa T. Psychosocial predictors of successful delivery after unexplained recurrent spontaneous abortions: a cohort study. *Acta Psychiatr Scand*. 2004;109:440-6.
17. Kim H, Jung Y. Effects of antenatal depression and antenatal characteristics of pregnant women on birth outcomes: a prospective cohort study. *J Korean Acad Nurs*. 2012;42:477.
18. Field T, Diego M, Hernandez-Reif M. Prenatal depression effects and interventions: a review. *Infant Behav Dev*. 2010;33:409-18.
19. Grote N, Bridge J, Gavin A, Melville J, Iyengar S, Katon W. A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Arch Gen Psychiatry*. 2010;67:1012-24.
20. Goodman SH, Gotlib IH. Risk for psychopathology in the children of depressed mothers: a developmental model for understanding mechanisms of transmission. *Psychol Rev*. 1999;106:458-90.
21. O'Connor T, Heron J, Glover V. Antenatal anxiety predicts child behavioral/emotional problems independently of postnatal depression. *J Am Acad Child Adolesc Psychiatry*. 2002;41:1470-7.
22. Plant D, Barker E, Waters C, Pawlby S, Pariante C. Intergenerational transmission of maltreatment and psychopathology: the role of antenatal depression. *Psychol Med*. 2012;43:519-28.
23. Deave T, Heron J, Evans J, Emond A. The impact of maternal depression in pregnancy on early child development. *Int J Obstet Gynaecol*. 2008;115:1043-51.
24. Murray L, Arteche A, Fearon P, Halligan S, Croudace T, Cooper P. The effects of maternal depression and child sex on academic performance at 16 years: a developmental approach. *J Child Psychol Psychiatry*. 2010;51:1150-9.
25. Rahman A, Lovel H, Bunn J, Iqbal Z, Harrington R. Mothers' mental health and infant growth: a case-control study from Rawalpindi, Pakistan. *Child Care Health Dev*. 2004;30:21-7.
26. Appleyard K, Berlin L, Rosanbalm K, Dodge K. Preventing early child maltreatment: implications from a longitudinal study of maternal abuse history, substance use problems, and offspring victimization. *Prev Sci*. 2011;12:139-49.
27. Urdaneta J, Rivera A, García J, Guerra M, Baabel N, Contreras A. Prevalencia de depresión posparto en primigestas y múltiples valoradas por la escala de Edimburgo. *Rev Chil Obstet Ginecol*. 2010;75:312-20.
28. Lam N, Contreras H, Mori E, et al. Factores psicosociales y depresión antenatal en mujeres gestantes. Estudio multicéntrico en tres hospitales de Lima, Peru: Abril a junio de 2008. *Rev Peru Epidemiol*. 2010;14.
29. Luna Matos ML, Salinas Piélago J, Luna Figueroa A. Depresión mayor en embarazadas atendidas en el Instituto Nacional Materno Perinatal de Lima. Peru. *Rev Panam Salud Pública*. 2009;26:310-4.
30. Bonilla-Sepulveda O. Depression and associated factors in pregnant and non-pregnant adolescent women in Medellín, Colombia, 2009. Cross sectional study. *Rev Colomb Obstet Ginecol*. 2009;2009.
31. Sharp D, Hay D, Pawlby S, Schmücker G, Allen H, Kumar R. The impact of postnatal depression on boys' intellectual development. *J Child Psychol Psychiatry*. 1995;36:1315-36.
32. Howard L, Oram S, Galley H, Trevillion K, Feder G. Domestic violence and perinatal mental disorders: a systematic review and meta-analysis. *PLoS Med*. 2013;10:e1001452.
33. Plant D, Barker E, Pariante C, Pawlby S. Multigenerational transmission of childhood maltreatment and psychopathology: the influence of antenatal depression. *Comprehens Psychiatry*. 2013;54:e8.
34. Berlin L, Whiteside-Mansell L, Roggman L, Green B, Robinson J, Spieker S. Testing maternal depression and attachment style as moderators of early head start's effects on parenting. *Attach Hum Dev*. 2011;13:49-67.
35. Fisher J, Cabral de Mello M, Patel V, Rahman A, Thach Tran T, Holtana S, et al. Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bull World Health Organ*. 2011;90:139-49H.
36. Sidebotham P, Heron J. Child maltreatment in the children of the nineties: a cohort study of risk factors. *Child Abuse Neglect*. 2006;30:497-522.
37. Beck C. A meta-analysis of predictors of postpartum depression. *Nurs Res*. 1996;45:297-303.
38. Robertson E, Grace S, Wallington T, Stewart D. Antenatal risk factors for postpartum depression: a synthesis of recent literature. *Gen Hosp Psychiatry*. 2004;26:289-95.
39. Boyce P. Risk factors for postnatal depression: a review and risk factors in Australian populations. *Arch Women Mental Health*. 2003;6:s43-50.

-
40. Rich-Edwards J. Sociodemographic predictors of antenatal and postpartum depressive symptoms among women in a medical group practice. *J Epidemiol Community Health*. 2006;60:221-7.
 41. Campo-Arias A, Ayola-Castillo A, Peinado-Valencia HM, Amor-Parra M, Cogollo Z. Validating the Edinburgh Postnatal Depression Scale's internal consistency and factor structure amongst pregnant women in Cartagena, Colombia. *Rev Colomb Obstet Ginecol*. 2007;58:277-83.
 42. Jadresic E, Araya R. Prevalence of postpartum depression and associated factors in Santiago de Chile. *Rev Med Chil*. 1995;123:694-9.
 43. Salazar-Pousada D, Arroyo D, Hidalgo L, Perez-Lopez F, Chedraui P. Depressive symptoms and resilience among pregnant adolescents: a case-control study. *Obstet Gynecol Int*. 2010;2010:1-7.