

# MEDICINA CLINICA PRÁCTICA



www.elsevier.es/medicinaclinicapractica

Images in medicine

# Cysticercosis - An unexpected finding

Cisticercosis - Un hallazgo inesperado

Susana Conceição\*,¹, Diogo Faustino¹, José Ribeiro²

Centro Hospitalar Universitário de Lisboa Central, Hospital de São José, Servico de Medicina 1.2, Portugal



## **Case description**

An 85-year old women with diabetes mellitus and a history of dementia was admitted to our medical ward with *C. difficile* colitis, following a previous episode of hospital admission due to a femoral fracture where she underwent antibiotic therapy to treat an infection of the resulting surgical wound. During this episode, abdominal and lower limb radiographs were taken (Fig. A and B), revealing several oval-shaped calcifications within the soft tissues and along the muscle fibres, aspects which are strongly suggestive of musculoskeletal Cysticercosis.

There was an accompanying history of abrupt, transient, visual impairment which, together with the aforementioned dementia, prompted an investigation to exclude neurological involvement (not confirmed). We found no larvae in the stools either. The patient passed away due to worsening of her infectious condition and no further testing was conducted.

Cysticercosis is a zoonosis caused by *Taenia solium* through ingestion of contaminated food products. Estimates point to an incidence of about twenty million new cases per year and around fifty thousand deaths. Embryo from ingested *T. solium* eggs gain access to the blood stream through




Fig. A. (Abdominal radiograph) and Figure B (Right Leg radiograph), both showing oval-shaped calcifications (cysticercus;).

<sup>\*</sup> Corresponding author at: Rua Cesário Verde, n° 28, Alto Estanqueiro/Jardia, 2870-620 Montijo, Portugal.

E-mail address: susana\_conceicao13@hotmail.com (S. Conceição).

<sup>&</sup>lt;sup>1</sup> Internal Medicine Specific Training

<sup>&</sup>lt;sup>2</sup> Senior Graduate Hospital Assistant of Internal Medicine

the hosts' intestinal mucosa and disseminate preferentially, through hematogenous spread, to nervous and muscle tissue. This distribution is responsible for the clinical features of the disease, although most of the infections (>80%) are asymptomatic.

This case highlights the typical radiographic aspects of musculoskeletal cysticercosis, which should prompt additional testing if found.

# **Ethical considerations**

The institution's protocols and procedures were followed regarding consultation and publication of patient data. Written consent was obtained from the patient's family for its publication, and their privacy was respected.

## **Funding**

No funding sources were received.

## **Conflict of interest**

The authors do not declare any conflict of interest.