




CIRUGÍA ESPAÑOLA

www.elsevier.es/cirugia



Video of the month

Intestinal cavernous hemangioma with a minimally invasive approach



Hemangioma cavernoso de intestino delgado con abordaje mínimamente invasivo

Antonio Enrique Gómez-Martín^{*}, Beatriz Martín-Pérez, Isabel María Gallarín-Salamanca, María Carmona-Agúndez

Servicio de Cirugía General y del Aparato Digestivo, Hospital Universitario de Badajoz, Spain

Cavernous hemangiomas represent between 5 and 10% of all benign tumors of the small intestine. Malignant transformation is rare and treatment is based on surgical or endoscopic resection. We present the case of a 26-year-old female with a 5-year history of iron deficiency anemia with initially negative work-up. Magnetic resonance enterography and capsule endoscopy revealed findings suggestive of hemangioma. An exploratory laparoscopy is carried out, locating an ileal hypervascular lesion, performing a laparoscopic intestinal resection and mechanic side-to-side anastomosis. Final pathology informed of a benign vascular lesion, compatible with cavernous hemangioma.

Conflict of interests

None.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.ciresp.2023.05.004](https://doi.org/10.1016/j.ciresp.2023.05.004).

Financing

None.

^{*} Corresponding author.

E-mail address: antonioenrique.gomez@salud-juntaex.es (A.E. Gómez-Martín).
<https://doi.org/10.1016/j.ciresp.2023.05.004>

0009-739X/© 2023 AEC. Published by Elsevier España, S.L.U. All rights reserved.