



Image of the month

Septic Shock Secondary to Mirizzi Syndrome Resolved in Two Stages[☆]



Shock séptico secundario a síndrome de Mirizzi resuelto en 2 tiempos

Adrián Huergo Fernández,^{*} Pedro Amor Martín, Marta Álvarez Posadilla

Servicio de Aparato Digestivo, Hospital Álvarez Buylla, Mieres del Camino, Asturias, Spain



Fig. 1




Fig. 2

A 70-year-old woman was admitted for acute cholecystitis and choledocholithiasis with clinical deterioration after 12 h due to septic shock secondary to cholangitis. Urgent ERCP detected dilatation of the proximal bile duct secondary to compression by a 2 cm calculus lodged in the cystic duct (Fig. 1). Given the patient's clinical situation, a double pigtail plastic stent was implanted to alleviate the biliary obstruction (Fig. 2), performing partial cholecystectomy and closure over the Kehr tube with extraction of the lithiasis 11 days later after the patient's status had improved.

[☆] Please cite this article as: Huergo Fernández A, Amor Martín P, Álvarez Posadilla M. Shock séptico secundario a síndrome de Mirizzi resuelto en 2 tiempos. Cir Esp. 2018;96:653.

^{*} Corresponding author.

E-mail address: adrianhuergo1978@gmail.com (A. Huergo Fernández).