



# Revista Española de Anestesiología y Reanimación

[www.elsevier.es/redar](http://www.elsevier.es/redar)



## CONSENSUS STATEMENT

### Spanish Society of Anesthesiology, Reanimation and Pain Therapy (SEDAR) Spanish Society of Emergency and Emergency Medicine (SEMES) and Spanish Society of Otolaryngology, Head and Neck Surgery (SEORL-CCC) Guideline for difficult airway management. Part II

M.Á. Gómez-Ríos<sup>a,\*</sup>, J.A. Sastre<sup>b</sup>, X. Onrubia-Fuertes<sup>c</sup>, Teresa López<sup>b</sup>,  
A. Abad-Gurumeta<sup>d</sup>, R. Casans-Francés<sup>e</sup>, D. Gómez-Ríos<sup>f</sup>, J.C. Garzón<sup>b</sup>,  
V. Martínez-Pons<sup>g</sup>, M. Casalderrey-Rivas<sup>h</sup>, M.Á. Fernández-Vaquero<sup>i</sup>,  
E. Martínez-Hurtado<sup>d</sup>, R. Martín-Larrauri<sup>j</sup>, L. Reviriego-Agudo<sup>k</sup>, U. Gutierrez-Couto<sup>l</sup>,  
J. García-Fernández<sup>m,n</sup>, A. Serrano-Moraza<sup>o</sup>, L.J. Rodríguez Martín<sup>p</sup>, C. Camacho Leis<sup>p</sup>,  
S. Espinosa Ramírez<sup>o</sup>, J.M. Fandiño Orgeira<sup>q</sup>, M.J. Vázquez Lima<sup>r,s</sup>, M. Mayo-Yáñez<sup>t</sup>,  
P. Parente-Arias<sup>t</sup>, J.A. Sistiaga-Suárez<sup>u</sup>, M. Bernal-Sprekelsen<sup>v,w</sup>, P. Charco-Mora<sup>g</sup>

<sup>a</sup> Anesthesiology and Perioperative Medicine, Complejo Hospitalario Universitario de A Coruña, A Coruña, Spain

<sup>b</sup> Anesthesiology and Perioperative Medicine, Complejo Asistencial Universitario de Salamanca, Salamanca, Spain

<sup>c</sup> Department of Anesthesiology, Hospital Dr Peset, Valencia, Spain

<sup>d</sup> Department of Anesthesiology, Hospital Universitario Infanta Leonor, Madrid, Spain

<sup>e</sup> Department of Anesthesiology, Hospital Universitario Infanta Elena, Valdemoro, Madrid, Spain

<sup>f</sup> Hospital de Barbanza, A Coruña, Spain

<sup>g</sup> Department of Anesthesiology, Hospital Universitari i Politècnic La Fe, Valencia, Spain

<sup>h</sup> Department of Anesthesiology, Complejo Hospitalario Universitario de Ourense, Ourense, Spain

<sup>i</sup> Department of Anesthesiology, Hospital Clínica Universitaria de Navarra, Madrid, Spain

<sup>j</sup> Department of Anesthesiology, Hospital Infanta Elena, Málaga, Spain

<sup>k</sup> Department of Anesthesiology, Hospital Clínico Universitario, Valencia, Spain

<sup>l</sup> Biblioteca, Complejo Hospitalario Universitario de Ferrol (CHUF), Ferrol, A Coruña, Spain

<sup>m</sup> Department of Anesthesiology, Hospital Universitario Puerta de Hierro-Majadahonda, Majadahonda, Spain

<sup>n</sup> President of the Spanish Society of Anesthesiology, Resuscitation and Pain Therapy (SEDAR), Spain

<sup>o</sup> SUMMA 112

<sup>p</sup> SAMUR Protección Civil, Madrid, Spain

<sup>q</sup> Servicio de Urgencias, Complejo Hospitalario Universitario de A Coruña, A Coruña, Spain

<sup>r</sup> Emergency Department, Hospital do Salnes, Vilagarcía de Arousa, Pontevedra, Spain

<sup>s</sup> President of the Spanish Emergency Medicine Society (SEMES), Spain

<sup>t</sup> Department of Otorhinolaryngology/Head Neck Surgery, Complejo Hospitalario Universitario A Coruña, A Coruña, Spain

\* Corresponding author.

E-mail address: [magoris@hotmail.com](mailto:magoris@hotmail.com) (M.Á. Gómez-Ríos).

<https://doi.org/10.1016/j.redare.2024.02.002>

2341-1929/© 2023 Sociedad Española de Anestesiología, Reanimación y Terapéutica del Dolor. Published by Elsevier España, S.L.U. All rights reserved.

Please cite this article as: M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al., Spanish Society of Anesthesiology, Reanimation and Pain Therapy (SEDAR) Spanish Society of Emergency and Emergency Medicine (SEMES) and Spanish Society of Otolaryngology, Head and Neck Surgery (SEORL-CCC) Guideline for difficult airway management. Part II, Revista Española de Anestesiología y Reanimación, <https://doi.org/10.1016/j.redare.2024.02.002>

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

<sup>u</sup> Department of Otorhinolaryngology, Hospital Universitario Donostia, Donostia, Gipuzkoa, Spain

<sup>v</sup> Department of Otorhinolaryngology, Hospital Clinic Barcelona, Barcelona Spain

<sup>w</sup> President of the Spanish Society for Otorhinolaryngology Head & Neck Surgery (SEORL-CCC)

Received 28 July 2023; accepted 28 August 2023

## KEYWORDS

Airway management;  
Practice guideline;  
Conscious sedation;  
General anaesthesia;  
Endotracheal  
intubation;  
Laryngeal mask;  
Tracheostomy;  
Airway obstruction;  
Monitoring;  
Rapid sequence  
induction;  
Airway extubation;  
Teaching

## PALABRAS CLAVE

Manejo de la vía  
áerea;  
Guía clínica;  
Sedación consciente;  
Anestesia general;  
Intubación  
endotraqueal;  
Mascarilla laríngea;  
Traqueostomía;  
Obstrucción de la vía  
área;  
Monitorización

**Abstract:** The Airway Management section of the Spanish Society of Anesthesiology, Resuscitation, and Pain Therapy (SEDAR), the Spanish Society of Emergency Medicine (SEMES), and the Spanish Society of Otorhinolaryngology and Head and Neck Surgery (SEORL-CCC) present the Guide for the comprehensive management of difficult airway in adult patients. Its principles are focused on the human factor, cognitive processes for decision-making in critical situations, and optimization in the progression of strategies application to preserve adequate alveolar oxygenation in order to enhance safety and the quality of care. The document provides evidence-based recommendations, theoretical-educational tools, and implementation tools, mainly cognitive aids, applicable to airway management in the fields of anesthesiology, critical care, emergencies, and prehospital medicine. For this purpose, an extensive literature search was conducted following PRISMA-R guidelines and was analyzed using the GRADE methodology. Recommendations were formulated according to the GRADE methodology. Recommendations for sections with low-quality evidence were based on expert opinion through consensus reached via a Delphi questionnaire.

© 2023 Sociedad Española de Anestesiología, Reanimación y Terapéutica del Dolor. Published by Elsevier España, S.L.U. All rights reserved.

## Guía de la Sociedad Española De Anestesiología, Reanimación y Terapéutica del Dolor (SEDAR), Sociedad Española de Medicina de Urgencias y Emergencias (SEMES) y Sociedad Española de Otorrinolaringología y Cirugía de Cabeza y Cuello (SEORL-CCC) para el manejo de la vía aérea difícil

**Resumen** La sección de Vía Aérea de la Sociedad Española De Anestesiología, Reanimación y Terapéutica del Dolor (SEDAR), la Sociedad Española de Medicina de Urgencias y Emergencias (SEMES) y la Sociedad Española de Otorrinolaringología y Cirugía de Cabeza y Cuello (SEORL-CCC) presentan la Guía para el manejo integral de la vía aérea difícil en el paciente adulto. Sus principios están focalizados en el factor humano, los procesos cognitivos para la toma de decisiones en situaciones críticas y la optimización en la progresión de la aplicación de estrategias para preservar una adecuada oxigenación alveolar con el objeto de mejorar la seguridad y la calidad asistencial. El documento proporciona recomendaciones basadas en la evidencia científica actual, herramientas teórico-educativas y herramientas de implementación, fundamentalmente ayudas cognitivas, aplicables al tratamiento de la vía aérea en el campo de la anestesiología, cuidados críticos, urgencias y medicina prehospitalaria. Para ello se realizó una amplia búsqueda bibliográfica según las directrices PRISMA-R y se analizó utilizando la metodología GRADE. Las recomendaciones se formularon de acuerdo con la metodología GRADE. Las recomendaciones de aquellas secciones con evidencia de baja calidad se basaron en la opinión de expertos mediante consenso alcanzado a través de un cuestionario Delphi.

© 2023 Sociedad Española de Anestesiología, Reanimación y Terapéutica del Dolor. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.


## Known or anticipated difficult airway

Awake management is the preferred option for securing the known or anticipated difficult airway (DAW) (expert state-

ment [E.S.] 85.7%)<sup>1</sup> because it (1) preserves airway (AW) patency and spontaneous ventilation, increases respiratory reserve, and confers protection against aspiration by preserving laryngeal reflexes,<sup>1,2</sup> (2) allows a gradual transition



## Anticipated difficult airway



**Figure 1** Cognitive aid to facilitate decision-making in the management of anticipated difficult airway.  
TI: tracheal intubation; DAW: difficult airway.

to positive pressure ventilation (PPV) and a slow induction of general anaesthesia (GA) given the risk for haemodynamic collapse,<sup>3,4</sup> (3) facilitates the procedure, because it inhibits soft tissue collapse, dilates peritracheal structures, allows for better location of the glottis by preventing the larynx from taking a more anterior position, and allows visualisation of air bubbles in the case of distorted anatomy, (4) allows the patient to remain in a sitting position, and cooperate, and enables assessment of their neurological status, and (5) allows all treatment options to remain open and decisions to be made according to the findings.<sup>1,5-8</sup>

A known or anticipated DAW requires a team-based strategy with thorough multidisciplinary discussion of sequential plans (primary and alternative) to achieve oxygenation, ventilation, and protection against aspiration.<sup>1,7,9,10</sup>

Safe management of DAW secondary to obstructive disease requires consideration of respiratory status, cause, location, and degree of obstruction (greater or less than 50%) through clinical signs and symptoms, imaging, and fibronasolaryngoscopy (FNL).<sup>1,7,10,11</sup> The risks and benefits of each approach should be carefully considered and the decision agreed by the medical and surgical team.<sup>1</sup> Fig. 1 shows a cognitive decision aid for the treatment of an anticipated DAW. The lower plans act as rescue strategies in the event of failure of the upper plan, which should be selected as the primary plan. Awake AW management is recommended in all cases. (1) Supraglottic lesions causing mild obstruction that can be overcome with an endotracheal tube (ETT) allow tracheal intubation (TI), usually with fibrobronchoscopy (FOB). (2) Obstructive supraglottic lesions with stenosis greater than 50% (or with inspiratory stridor at rest 50%),<sup>7</sup> non-obstructive supraglottic lesions that prevent TI

(or associate TI with unacceptable morbidity), and glottic or subglottic lesions make tracheostomy or cricothyrotomy advisable as the primary approach.<sup>10</sup> (3) Lower tracheal obstructive lesions not salvageable with an ETT or tracheal cannula require extracorporeal membrane oxygenation (ECMO).<sup>10</sup>

Active oxygenation strategies should be implemented throughout the procedure. High-flow nasal oxygen therapy (HFNO), although it requires further validation in this setting, may be the technique of choice. HFNO is recommended over conventional low-flow nasal cannula (E.S. 91.4%).

### Awake tracheal intubation

When it is feasible to secure the AW non-invasively, awake TI is still the gold standard for DAW management<sup>3,12,13</sup> because it is safe and reliable.<sup>14,15</sup> Four elements are critical to the success of the procedure: continuous oxygenation, topicalization of AW, sedation (optional), and selection, experience, and appropriate management of the TI device and technique. The ideal protocol in terms of efficacy and safety is unknown, and therefore the most appropriate protocol must be chosen according to the clinical context and individual patient characteristics, and the operator's experience and preferences.<sup>16-18</sup> Fig. 2 shows the aid proposed by the Spanish Society of Anaesthesiology, Reanimation and Pain Therapy (SEDAR) Spanish Society of Emergency and Emergency Medicine (SEMES), and Spanish Society of Otolaryngology, Head and Neck Surgery (SEORL-CCC) for TI of the anticipated DAW.




# Anticipated difficult airway



## Position – Ergonomics

### Oxygenation + topicalisation ± sedation



**Figure 2** Cognitive aid proposed by SEDAR SEMES for tracheal intubation of unanticipated difficult airway.  
ATI: awake tracheal intubation. DAW: difficult airway; Int2: second operator.

## Oxygenation

Continuous oxygenation increases safety by preventing or minimising hypoxaemia.<sup>12,19</sup> Conventional methods may be insufficient to prevent desaturation.<sup>20</sup> HFNO allows longer tolerance of potential AW obstruction, hypoventilation, or apnoea period.<sup>20–23</sup> Therefore, although the evidence is incipient, it is becoming the method of choice.<sup>13,24</sup>

Non-invasive ventilation (NIV) with endoscopic mask may have a role in IT of critically ill patients with severe hypoxaemia (E.S. 82.9%).<sup>25</sup>

## Topicalisation

Topical anaesthesia of the AW is the key element of the procedure, because it is the main determinant of its success.<sup>1</sup> With its good cardiovascular and systemic toxicity risk profile, lidocaine (2%–4%) is the most widely used local anaesthetic (LA).<sup>12,19,26–28</sup> The maximum total dose should not exceed 9 mg/kg.<sup>13,29</sup> The minimum dose necessary should be used.

The spray-as-you-go (SAYGO) technique, using an epidural catheter or atomisers, and regional blocks (glossopharyn-

geal, superior laryngeal, transtracheal injection) are the most commonly used methods for topical anaesthesia of the respiratory mucosa.<sup>2,12,27</sup> They are usually used in combination.<sup>28</sup> There is no evidence as to which method is superior,<sup>16</sup> although nerve blocks are more invasive, require multiple bilateral injections, require experience, and are associated with a higher incidence of complications.<sup>2,13,16</sup> Transtracheal block (lidocaine 4%, 4 mL) is perhaps the most useful invasive method as (1) it provides anaesthesia of the subglottic larynx, upper trachea, and even supraglottic structures,<sup>30</sup> (2) its success rate is greater than 95%,<sup>12</sup> and (3) complications are rare (1:10,000), although they can be significant.<sup>28,31</sup>

The SAYGO technique can minimise the risk of aspiration, as laryngeal reflexes are maintained until just before passing the ETT.<sup>2</sup> Premedication with an antisialogogue is recommended to optimise LA efficacy and field of view, glycopyrrrolate (3 µg/kg) being the antisialogogue of choice (E.S. 80%) due to its rapid onset of action, nil effect on the central nervous system (CNS), and moderate vagolytic effect.<sup>2,12,32</sup> Its administration 15–20 min beforehand reduces the dilution and oesophageal elimination of LA by secretions.<sup>1</sup>

Regardless of the method used, topicalisation should include the oral cavity, nasal cavity if nasotracheal intubation is planned, oropharynx, periglottic area, larynx, and trachea.<sup>1,30</sup> Otherwise, device insertion and TI trigger reflex responses of the AW, such as cough or laryngospasm, as well as a cardiovascular response mediated by the sympathetic nervous system.<sup>33</sup>

## Sedation

Sedation is an optional adjunct to adequate topical anaesthesia in awake airway management (ATI) (E.S. 88.6%)<sup>16</sup> because awake TI with prior psychological preparation can be safely and effectively performed without it.<sup>19,28,34</sup> It should never compensate for poor topicalisation. While very high levels of anxiety may increase the physiological stress response and reduce tolerance, over-sedation may cause loss of cooperation, respiratory depression, hypoxia, hypercapnia, AW obstruction, aspiration, or cardiovascular instability.<sup>19,35,36</sup> Therefore, the administration of sedation must be justified through a thorough risk-benefit analysis. The objectives to be achieved are (E.S. 94.3%),<sup>1,19,35</sup> (1) effective anxiolysis and amnesia while maintaining patient cooperation (conscious sedation; Ramsay sedation score 2–3)<sup>28,37</sup> and (2) analgesia to suppress cough and gag reflex, and reduce haemodynamic response while preserving AW patency, spontaneous ventilation, and avoiding aspiration. Careful monitoring is required to avoid overdosing and it is recommended that a second anaesthetist be exclusively responsible for injecting and monitoring the effects of sedation.<sup>13,19</sup>

The different sedation regimens have shown a satisfactory level of efficacy and safety.<sup>16</sup> Dexmedetomidine, with its anxiolytic, sedative, analgesic, and sympatholytic properties, may have a better efficacy and safety profile by producing fewer episodes of apnoea and desaturation, and is well tolerated and associated with better TI conditions and lower recall compared to other drugs,<sup>16,38–40</sup> although it may cause episodes of severe bradycardia and hypotension.<sup>35</sup> Its ability to maintain the integrity of respiratory function, even with deep levels of sedation, makes it a good choice for patients at risk of AW obstruction and/or respiratory failure.<sup>35</sup> Opioids, particularly remifentanil, attenuate cough and gag reflex, although they may increase the incidence of chest rigidity and laryngospasm<sup>6,13,19</sup> and are linked to a high incidence of recall when used as a single drug, requiring administration with a benzodiazepine such as midazolam.<sup>19,28</sup> In general, monotherapy is more predictable and reliable, although its safety is enhanced with the availability of specific opioid and benzodiazepine antagonists. Remifentanil is a good option when topical anaesthesia is not possible.<sup>35,41</sup> Propofol is associated with a low incidence of recall at the expense of an increased risk of excessive sedation, AW obstruction, and cough.<sup>13,35,42</sup> Table 1 shows the main drugs used for sedation.

## Device

FOB-guided TI is classically considered the method of choice in the awake patient<sup>2,17,37,43</sup> because of its versatility and unique combinability with other devices in any treatment

plan,<sup>44</sup> and its efficacy and safety.<sup>16,45</sup> However, it is a complex skill that requires regular practice, is fallible, and is not available for all patients.<sup>37,46</sup>

Elective awake TI with video laryngoscopy (VL) may be faster than FOB-guided, which may decrease the risk of aspiration, and be associated with equivalent success rates, safety profile, and patient and operator satisfaction.<sup>17,43,47,48</sup> Furthermore, the competence required for its use is easy to acquire and maintain.<sup>46,47</sup> It is therefore a valid alternative technique as a first-line treatment.<sup>13,17,46–49</sup> The video laryngoscopes indicated for this context are those with a hyperangulated blade, with or without a guide channel. In certain circumstances, VL offers additional advantages over FOB.<sup>17,37,43</sup> It also allows an ETT of any diameter to be selected,<sup>37</sup> it can be changed without removing the device,<sup>47</sup> and its passage through the vocal cords can be observed, unlike advancing blind with FOB, which reduces the risk of injury from the ETT and associated trauma.<sup>37,50,51</sup> However, unlike FOB, its use is not feasible with a buccal opening of less than 18–20 mm or a space-occupying lesion in the oral cavity,<sup>37</sup> and may result in increased cervical spine movement in patients with instability without manual stabilisation,<sup>52</sup> although the results appear similar.<sup>53,54</sup> Therefore, VL cannot completely replace FOB-guided TI.<sup>37,43</sup>

There is insufficient evidence as to which technique is ideal, and therefore selection of the device must be context-sensitive.<sup>43,48</sup> The two approaches are virtually equivalent and complementary<sup>43</sup>; a failed attempt of one can be rescued with the alternative and both can be used in combination, primarily to treat very difficult AW.<sup>2,17,37</sup>

Single-use flexible FOBs have a similar safety profile to reusable FOBs,<sup>55,56</sup> although they may be associated with benefits in terms of cost-effectiveness, cross contamination, infection, and resource utilisation.<sup>57,58</sup>

## Procedure

The conditions of all four components should be optimal from the first attempt to maximise the likelihood of success and minimise the number of attempts. If the selected primary technique (FOB or VL) fails, the alternative technique should be used (E.S. 80%). A third attempt may benefit from a multimodality approach consisting of the combination of VL + FOB (E.S. 100%), therefore using the advantages of both devices.<sup>59</sup> This combination may improve the first attempt success rate, reduce TI time, reduce morbidity,<sup>60,61</sup> and could be the approach of choice in AWs with severely distorted anatomies.

The combination of a supraglottic airway device (SAD) and FOB may be useful as a rescue technique to maintain oxygenation, maintain AW patency, and perform TI through the SAD (E.S. 100%).<sup>62,63</sup> The SAD acts as a conduit to for an FOB-guided TI, maintaining oxygenation and AW patency and isolating the periglottic structures from any secretions or blood. At the same time, it facilitates locating the glottis and decreases the difficulty of advancing the ETT over the FOB,<sup>51</sup> while the latter allows the SAD to be placed in the correct location. The new laryngeal video masks could simplify this technique.<sup>65,66</sup>

The total number of attempts should be limited to three (E.S. 88.6%) because repeated attempts increase the risk

**Table 1** Main drugs used for sedation in awake airway management.

Drug	Dose	Effects	Advantages	Disadvantages
Dexmedetomidine	Bolus of de. –1.0 µg/kg over 10 min; followed by infusion of .5–1.0 µg/kg/hour	Sedation/analgesia/amnestic antisialogogue	Respiratory profile Safe	Bradycardia, hypotension
Remifentanil	Infusion of .03–.1 µg/kg/min; Bolus of .05–.1 µg/kg	Analgesia/antitussive	Suppresses cough reflex	Respiratory depression Laryngeal reflex depression Recall of procedure
Midazolam	Boli of .015–.03 mg/kg	Amnesia/sedation	Concomitant use with opioids Amnesia Minimises side effects	Respiratory depression
Fentanyl	Boli of .7–1.5 µg/kg	Analgesia/antitussive	Rapid onset	Respiratory depression Laryngeal reflex depression
Propofol	Infusion of 25–75 µg/kg/min; bolus of 25–75 µg/kg	Sedation	Amnesia Synergistic effect with other drugs (reduction in doses required)	Oversedation Airway obstruction Episodes of hypoxaemia Does not suppress cough reflex
Ketamine	Boli of .07–.15 mg/kg	Sedation/analgesia	Preserves muscle tone and airway protective reflexes Opioid sparing effect Concomitant use with dexmedetomidine increases haemodynamic stability	Agitation (co-administration with midazolam to avoid), inadequate level of sedation, intense cough, and unpleasant recall Increased secretions (requires antisialogogue as premedication) Myocardial depression in states of catecholamine depletion

Doses taken from Gil K, et al.<sup>6</sup>

of oedema, laryngeal bleeding, and complete obstruction of the AW.<sup>51</sup> It is especially important in the presence of a pre-existing obstruction because it can rapidly progress to a can't intubate-can't-oxygenate (CICO) situation.<sup>9,67</sup> Each failure should be followed by an assessment of each of the four components of the technique to determine whether optimisation is feasible.

Factors such as surgical approach, anatomical features, extubation plan, and operator preference determine the choice of TI route.<sup>68</sup>

The inability to advance the ETT over the FOB, stylet, or an exchanger through the vocal cords due to impingement upon periglottic structures, primarily the right arytenoid cartilage,<sup>2,50</sup> is one of the main reasons for TI failure,<sup>19,51</sup> and glottic or subglottic damage.<sup>50,69</sup> ETTs with a conical opening made of silicone such as LMA Fastrach™ (Teleflex Medical, Dublin, Ireland),<sup>70</sup> those with a central opening and posterior facing bevel such as the Parker Flex-Tip™ (Parker Medical, Highlands Ranch, CO USA),<sup>71–74</sup> LMA Fastrach™,<sup>75</sup> and BlockBuster (Tuoren Medical Instrument co, Ltd, Changyuan city, China),<sup>69</sup> and flexible material and reduced or no anterior curvature such as the reinforced

LMA Fastrach™ and flexometallic ETTs<sup>69,76</sup> reduce the incidence of this complication.<sup>50,69,72,76,77</sup> The Parker Flex ETT is recommended over conventional ETT for FOB-guided TI in the general population (1 B), posterior placement of the bevel or 90° anti-clockwise rotation so that the bevel faces posteriorly facilitate its advancement.<sup>78–80</sup> Other useful manoeuvres include cervical spine flexion and release of jaw thrust or cricoid pressure.<sup>51,80</sup> It is recommended to decrease the difference between the external diameter of the FOB and the internal diameter of the ETT to facilitate FOB-guided TI (E.S. 85.7%),<sup>2,51,81</sup> or to use an intubation catheter between the FOB and the ETT (e.g. Aintree catheter, AIC, Cook Critical Care, Bloomington, IN, USA). Similarly, a device-specific ETT is recommended facilitate VL-guided TI.<sup>82</sup> Parker Flex ETT rather than conventional ETT is suggested for FOB and laryngoscopy in the general population to reduce complications (1 C).

Smaller sized ETTs allow a better laryngeal view during laryngoscopy and facilitate TI by reducing the impact on periglottic structures.<sup>51,83,84</sup> Also, large ETTs are associated with increased morbidity.<sup>83,85–90</sup> ETTs up to 6.0 mm internal diameter allow access for intubation devices, suc-

tion devices, and small-calibre FOB. PPV can be performed without increasing the risk of ventilator-induced lung injury or air trapping,<sup>91,92</sup> even when high minute volumes (MV) are required. There is also no demonstrable increased risk of aspiration or cuff pressure damage when smaller ETTs are used; they may even provide a better seal than the larger ETTs.<sup>93</sup> Smaller ETTs may not be safe in all cases, such as in patients with heavy secretions or limited airflow. Therefore, a smaller ETT than usual is recommended with VL and FOB (D.E. 85.7%) for safe treatment performing a risk-benefit analysis according to the clinical context.<sup>83,84,90,94</sup>

After visual confirmation of TI (passage of the ETT through the glottis with VL and identification of the carina and advancement of the ETT over FOB to two to three tracheal rings above the carina) it is recommended to induce GA after establishing cuff pressure and capnographic confirmation of TI (E.S. 94.3%).<sup>2</sup>

Alternative techniques and approaches should be planned in advance and applied without delay after failure of the primary approaches (E.S. 100%).<sup>12,67</sup> In the event of failed awake TI, there are three alternative plans: (1) postponement of the procedure unless the situation requires immediate treatment, (2) awake front-of-neck access (FONA), or emergency FONA in case of obstruction and loss of AW control, (3) GA induction with complete neuromuscular blockade (NMB) and follow-up of the unexpected DAW algorithm. This is a high-risk option. Deep sedation or GA with spontaneous ventilation induced with ketamine might be a more favourable preliminary step than the establishment of apnoea<sup>95</sup> (Fig. 2).

Given the high likelihood of awake TI failure, it is recommended to prepare the FONA plan in parallel (double configuration) to treat a potential total obstruction (E.S. 88.6%)<sup>96</sup>; the incision line is anticipated by marking the cricothyroid membrane (CTM),<sup>3</sup> the neck and material are prepared with the surgical team present.<sup>9</sup> Multidisciplinary management and coordination with ENT is essential.<sup>1,97</sup>

## Recommendation

A Parker Flex ETT is recommended over conventional ETT for FOB-guided TI in the general population.

**Strong recommendation; moderate level of evidence** (⊕⊕⊕⊖)

A Parker Flex ETT is recommended over conventional ETT for FOB-guided TI in the general population.

**Strong recommendation; low level of evidence** (⊕⊕⊖⊖)

## Alternative awake techniques

### Front-of-neck access technique

Awake tracheostomy under local anaesthesia is recommended in the presence of pre-existing critical AW compromise (E.S. 82.9%). In the upper AW with significant distortion or obstruction due to neoplasia, haematoma,

severe oedema, bilateral vocal cord paralysis, or haemorrhage, awake FONA under local anaesthesia as the primary plan may be the safest option to secure the AW,<sup>67,98–102</sup> because, (1) instrumentation of the upper AW may precipitate bleeding, oedema, increased occlusion, and even distal seeding of a tumour,<sup>7,9</sup> (2) topical anaesthesia may exacerbate a pre-existing occlusion,<sup>1,103</sup> or cause laryngospasm,<sup>1,9</sup> and (3) FOB can cause a cork in a bottle effect<sup>7</sup> with complete collapse. Multidisciplinary DAW teams are associated with better first attempt success rates and faster time to securing a DAW.<sup>104</sup>

The recommended technique is tracheostomy performed by a trained professional.<sup>105</sup> Sedation should be avoided if possible. HFNO appears to be effective in extending safe apnoea time,<sup>106</sup> although extreme caution should be exercised and the use of electrical instruments should be limited at the time of tracheal opening.<sup>107</sup> The procedure requires the patient's cooperation as supine decubitus and cervical spine extension is often poorly tolerated.<sup>106</sup> Once a capnographic record is available, anaesthetic induction is performed.<sup>108</sup> The full moon view of the wall of trachea with FOB confirms correct cannula location. A crescent image indicates the need to reposition or change the cannula.

Awake cricothyrotomy would be the most indicated technique for emergent critical airway compromise (D.E. 91,4%) as it enables a surgical AW to be established more rapidly.<sup>109</sup>

## Extracorporeal membrane oxygenation

ECMO under local anaesthesia in the awake patient may be the safest option when all four conventional plans are expected to be impossible, unsuccessful, or ineffective with risk of complete obstruction of the AW (E.S. 90.6%). Technological advances have made it possible to incorporate ECMO in AW management in these situations to ensure adequate oxygenation.<sup>110,111</sup> This is the case of tracheobronchial or extrinsic pathologies, such as anterior cervical spine pathology, which result in a critical central airway obstruction or prevent the performance of an FONA.<sup>67,110</sup> In these cases, elective ECMO under local anaesthesia in the awake patient may be the safest option.<sup>67,112,113</sup> Once a controlled environment is achieved, the AW is secured to prevent aspiration. Table 2 shows the main indications for ECMO.

Given the cost and its potential complications,<sup>111,114–116</sup> ECMO must be supported by a careful multidisciplinary decision-making process.<sup>116</sup> In uncertain indications, a system prepared with cannulated vessels may be possible and a perfusionist present prior to AW management.<sup>116</sup>

The respiratory support provided by veno-venous ECMO is associated with fewer complications, does not require therapeutic levels of anticoagulation, and allows the use of a single double-lumen cannula, making it the technique of choice in these situations.<sup>110,111</sup> In cases with associated haemodynamic compromise requiring cardiorespiratory support, such as large mediastinal masses, veno-arterial ECMO, or even a conventional cardiopulmonary bypass circuit may be necessary.<sup>110</sup>

In extreme cases such as massive haemoptysis or central foreign bodies, ECMO may be a last resort.<sup>64,117</sup> However, setting it up can be complicated and requires considerable time, so it cannot currently be considered a rescue

**Table 2** Main indications for extracorporeal membrane oxygenation (ECMO).**Tracheobronchial pathology**

- Tracheobronchial tumours with critical stenosis
  - Tracheal stenosis of non-tumour cause (sequelae of prolonged intubation or tracheostomy, obstructive tracheal papillomatosis)
  - Tracheal deformities
  - Tracheal trauma (accidental or iatrogenic)
  - Haemoptysis (the need for anticoagulation may complicate haemostasis)
  - Pathologies requiring complex surgery (bronchopleural fistulae, tracheoesophageal fistulae, carinal resections)
  - Complications arising from tracheal stents
  - Foreign bodies in the airway
- Extrinsic pathology**
- Thyroid neoplasm or large goitres with severe tracheal invasion or compression
  - Mediastinal masses with severe compression of the airway and/or great vessels, or heart chambers

technique for a CICO situation after GA induction, although several published cases have described its use in this context.

**Unanticipated difficult airway****Peri-procedural oxygenation**

Covered in the relevant section.

**Airway management**

TI is associated with more complications than other non-invasive plans, and therefore it should not be a rash act; but an active decision and should be only performed when truly indicated.<sup>118</sup>

Any approach to the AW should aim to minimise the number of attempts to avoid a CICO situation and the need for FONA. Planning and optimisation play a key role.<sup>119</sup>

Given the unreliability of predictors,<sup>120,121</sup> planning should be geared towards managing a potentially difficult AW.<sup>122</sup> Decision-making should be context-sensitive, rather than focussed on specific devices and techniques.<sup>8,123</sup>

The first attempt should be under a priori optimal conditions (E.S. 100%) to maximise the probability of success (make your first attempt your best attempt).<sup>119,122,124–126</sup> Additional attempts are only justified when there is room for improvement and involve the summative effect of optimisation, or allow substantial improvement in the likelihood of success (e.g., change the size, type of device, adjuvant, or operator as needed).<sup>119,127</sup>

Between attempts, face mask ventilation (FMV), and anaesthetic depth and NMB should be checked. Adequate ventilation between attempts with face mask (FM) or second-generation SAD (2 GSAD) provides an opportunity to stop and think to reformulate the strategy, or deploy new resources while maintaining first principles. Fig. 3 shows the treatment algorithm for unexpected TI.

Failure of non-invasive plans requires declaration of a CICO situation, ensuring adequate NMB, and immediate FONA regardless of peripheral oxygen saturation value ( $\text{SpO}_2$ ).

**Tracheal intubation****Videolaryngoscopy**

Failure of the first attempt at TI implies a lower probability of success in subsequent attempts.<sup>60,119,128</sup> Multiple attempts may result in trauma, oesophageal intubation, hypoxaemia, cardiovascular events, a CICO situation, unexpected admission to the critical care unit, or death.<sup>129–134</sup> In addition, up to 93% of difficult TIs are unanticipated.<sup>121</sup> Therefore, the most appropriate primary technique should be the one that offers the highest guarantee of a first-attempt TI (E.S. 94.3%).<sup>119,122,124–126</sup>

Most meta-analyses, despite great heterogeneity<sup>135,136</sup> suggest the superiority of VL (Appendix A, supplementary material 4). Overall, VL compared to direct laryngoscopy (DL) increases success at the first attempt,<sup>137–150</sup> improves visualisation of the glottis,<sup>140,141,143,145–148,150–158</sup> and decreases complications, primarily trauma, and the incidence of oesophageal intubation<sup>141,142,144–147,149,152,155–157,159–162</sup> by as much as 50%.<sup>150</sup>


The emergence of the COVID-19 pandemic,<sup>135,163–167</sup> the reduction in costs<sup>18</sup> and widespread availability, and incipient positive data on cost-effectiveness,<sup>168,169</sup> quality of care,<sup>129</sup> teaching, documentation, teamwork (it favours the shared mental and human factors [HF] model),<sup>122,167,170–173</sup> and mastery of the technique with regular practice<sup>129</sup> have overcome resistance to change,<sup>174,175</sup> making VL practically the gold standard for laryngoscopy and DAW.<sup>171</sup>

In view of the above, SEDAR SEMES SEORL-CCC recommends the routine use of VL over DL as the primary device for TI (1 B). Devices with standard Macintosh-type blades (allowing direct and indirect laryngoscopy) are appropriate for AW management without predictors of difficulty, while those with hyperangulated blades (with or without guide channel) are indicated for known or anticipated DAW (E.S. 94.3%).<sup>120,135,176,177</sup> The latter are the devices of choice for rescue after a first unsuccessful attempt.<sup>14,178–180</sup> It is therefore recommended that VL and the necessary competence be available in any location where the AW is managed.

Inability to pass the ETT through the glottis is the most common cause of failed TI with VL.<sup>166,172,173,181,182</sup> However, with adequate competence it is rare for this problem not



## Unanticipated difficult tracheal intubation



\* Each "Plan" or non-invasive strategy should be limited to a maximum of 3 attempts.

\* Continuous oxygen therapy throughout the procedure (FMV/VSAD between attempts and apnoeic oxygenation during them)

\* Between attempts, the level of anaesthetic depth and NMB should be checked, and whether there are possibilities for optimisation

**Figure 3** Treatment algorithm for unanticipated tracheal intubation.

† Ventilation grades according to capnography waveform; CRICO: cricothyrotomy; DAW: difficult airway; DL: direct laryngoscopy; FONA: front-of-neck access; ETT: endotracheal tube; FMV: face mask ventilation; SAD: supraglottic airway device; SpO<sub>2</sub>: peripheral oxygen saturation; TI: tracheal intubation; VSAD: ventilation with supraglottic airway device; VL: videolaryngoscopy.

\*Each "Plan" or non-invasive strategy should be limited to a maximum of 3 attempts.

\*Continuous oxygen therapy throughout the procedure (FMV/VSAD between attempts and apnoeic oxygenation during them).

\*Between attempts, the level of anaesthetic depth and NMB should be checked, and whether there are possibilities for optimization.

to be overcome.<sup>171</sup> The recommended manoeuvres to overcome this difficulty are shown in Table 3.<sup>182-186</sup>

Bougies are associated with higher first-attempt success rates in elective and emergency TI, especially in patients with predictors of DAW or poor view of the glottis.<sup>187-190</sup> It is recommended that a bougie be available in every location with AW management (E.S. 97.1%)<sup>191</sup> and its use should be assessed from the first attempt.<sup>120,185,188,192</sup> SEDAR SEMES SEORL-CCC recommend the use of a dynamic or articulated bougie versus a conventional stylet for TI in patients with DAW (1 C) because it improves the first-attempt TI rate and time to successful intubation, thus reducing instrumentation and the use of alternative adjuvants.<sup>60,69,186,193,194</sup>

## Recommendation

### Flexible fibrobronchoscopy

FOB-guided TI in the patient, unconscious or under GA can be very effective in skilled hands,<sup>195-198</sup> but is technically more difficult than in the awake patient, and it may fail and be associated with episodes of desaturation or complete obstruction of the AW.<sup>105</sup> The presence of blood, emesis, or secretions in emergency TI further reduces the likelihood of success.

**Table 3** Manoeuvres to overcome difficult tracheal intubation with video laryngoscopy.

**Relating to the relationship of the device to the glottis to optimise view**

- Selection of an appropriate device size, especially those with a guide channel. The internal size of the guide channel determines the orientation of the ETT exit.
  - Adjust the distal position of the device with partial elevation and retraction to enlarge the field of view and align the blade to the glottis. In the case of devices with a guide channel, this allows centring the view of the glottis and thus optimising the TI, as the ETT advances towards the glottis under a predetermined angle defined by the channel configuration and the angulation of the distal end of the ETT.
  - External laryngeal manipulation (BURP)
  - Increase head elevation
  - Clamp the epiglottis with the blade in case of large hanging epiglottis obstructing the view of the glottic structures and passage of the ETT

**Relative to the use of adjuvants**

- Semi-rigid stylet, either malleable, which can be pre-formed as a ‘hockey stick’ to adapt the ETT to the same angulation as the blade of the device, or the manufacturer’s own stylet designed for use with a specific VL. Their use is indispensable in the case of VLs with a hyper-angulated blade without a guide channel. As soon as the vocal cords are passed, it is recommended to remove the stylet to avoid injury to the AW.
- Static bougie with angled distal end. Limited usefulness if they are not malleable as they do not maintain curvature similar to that of the blade and for devices with a guide channel, more useful for video laryngoscopes with Macintosh-type blade.
- Dynamic bougie or combined use with flexible FOB to negotiate the acute angle between the glottis and the distal end of the ETT.

**Relative to the endotracheal tube**

- Type of ETT. Flexible tubes with silicone distal end and conical tip
  - Appropriate size of ETT. In general, a smaller size than usual is recommended to facilitate passage through the glottis, although excessively small ETTs in devices with a guide channel can hinder TI to the extent that by widening the difference between the internal calibre of the channel and the external calibre of the ETT, the exit of the distal end of the ETT is separated from the distal end of the blade, and takes an excessively posterior position.
  - Modify the curvature of the ETT
  - ETT with flexible articulating distal end that allows articulation at a variety of angles.
  - Rotating the ETT 90° clockwise so that the bevel faces forward reduces the angle of incidence of the ETT when intubation adjuvants are not used, or the stylet is removed prior to advancing the ETT into the trachea. When using a dynamic bougie or combined flexible FOB, rotating the ETT 90° is recommended so that the bevel is posterior facing to overcome the difficulty in advancing it.

BURP: backward, upward, right lateral position; FOB: fibrobronchoscopy; TI: tracheal intubation; ETT: endotracheal tube; AW: airway; VL: videolaryngoscopy.

Routine use of VL is recommended over DL as the primary device for TI.

**Strong recommendation; moderate level of evidence** (⊕⊕⊕⊖)

A dynamic or articulated bougie (flex-tip or FOB type) is recommended over a conventional stylet for TI in patients with DAW.

**Strong recommendation; low level of evidence** (⊕⊕⊖⊖)

Manoeuvres such as lingual traction and jaw-thrust allow opening of the pharynx and larynx respectively, improving view and the success of the technique.<sup>197,199</sup> Oral canulae, anterior displacement of the base of the tongue with the laryngoscope or patient in the left semilateral position rotating their head to the left facilitate passage of the FOB and improve view.<sup>200</sup> If there is resistance to ETT advancement, jaw-thrust and anticlockwise rotation of the ETT can facilitate its passage through the glottis.<sup>198</sup>

All the patients included in the 4th National Audit Project (NAP4) registry in whom FOB-guided TI was attempted after induction of GA as the primary technique or after failure of DL, failed and emergency FONA was required.

FOB is highly salvageable after failure of most devices, either alone or as a multimodal approach.<sup>6</sup> In an emergency, its availability, preparation, and execution is more laborious and time-consuming than VL,<sup>43,47</sup> and therefore it is less widely used as a rescue device.<sup>201</sup>

### Confirmation of tracheal intubation

Once TI has been achieved, it is recommended to immediately rule out oesophageal intubation as this is a common complication with devastating consequences.<sup>167,202–204</sup>

No confirmatory technique is 100% reliable in all circumstances, and therefore a combination of methods is recommended.<sup>205,206</sup>

The capnography waveform is the gold standard for confirming TI (1 B) because it is the most accurate (sensitivity 98%–100% and specificity 100%) and rapid method.<sup>4,120,205,207–212</sup> Therefore, despite persistent deficits

in standardisation,<sup>204</sup> capnography is mandatory to confirm ETT placement and should be present in all locations where AW management may be required.<sup>167,203,211,213-216</sup> An absent trace (grade 3 ventilation) indicates failed TI until proven otherwise (E.D. 80%)<sup>211,217</sup> and oesophageal intubation should be actively excluded.<sup>218</sup> The capnography trace is present but attenuated (not flat) even in cardiac arrest.<sup>211,212,217-219</sup> The positive predictive value of end-tidal carbon dioxide (EtCO<sub>2</sub>) is lower during cases of low or no perfusion<sup>212</sup> and false positive readings are possible when the ETT tip is in the hypopharynx.<sup>205</sup> In these cases, confirmation by ultrasound is a valid alternative as it has a sensitivity of 99% and specificity of 97%, is independent of pulmonary blood flow, is quick to perform (mean time of 13 seconds), and even allows real-time visualisation of the introduction of the ETT into the trachea or oesophagus.<sup>220,221</sup> Presence of the double tract sign allows detection of oesophageal intubation before initiation of ventilation.<sup>216,222</sup>

Colorimetric capnometry should be limited to situations where a capnography waveform is not available, such as in pre-hospital or emergency settings.<sup>205</sup>

Clinical examination alone has too high a false positive rate,<sup>223</sup> especially in an emergency setting,<sup>224</sup> and confirmation bias can lead to seeing and hearing what one wants,<sup>225</sup> although in combination with capnography it is useful. Ultrasound or fibrobronchoscopy examination via the ETT are alternative methods of confirming a TI. Other methods are oesophageal detector devices, transtracheal illumination, pulse oximetry, and chest radiography.<sup>205</sup>

Capnography waveform monitoring during maintenance of mechanical ventilation is highly recommended in all settings (E.S. 100%)<sup>207,209,211,226-230</sup> because it allows continuous monitoring of the position of the ETT, confirmation of AW patency, and early diagnosis of inadvertent extubation or partial displacement of an artificial AW.<sup>211,218,227,229,231-233</sup> The NAP4 registry found that lack of monitoring may have contributed to more than 70% of AW-related ICU deaths,<sup>234</sup> and therefore widespread use of capnography in critical care units and education of medical and nursing staff<sup>211,217,218</sup> may be the single change with the greatest potential to prevent morbidity and mortality associated with TI or other artificial AW outside the operating theatre.<sup>211,234</sup>

## Recommendation

Capnography waveform is recommended as the gold standard to confirm alveolar ventilation.

**Strong recommendation; moderate level of evidence** (⊕⊕⊕⊖)

## Ventilation with supraglottic airway device

In addition to their use as a primary technique in elective surgical procedures or cardiopulmonary resuscitation,<sup>235-237</sup> SADS play an indispensable role in the rescue of difficult or failed TI as they allow ventilation and oxygenation, provide

a patent AW with some degree of protection against aspiration, and act as conduits to facilitate FOB-guided TI.<sup>238-243</sup> Anatomical and/or technical factors hindering FMV and IT usually do not influence the insertion and function of the SAD.<sup>239</sup> Therefore, an SAD should be inserted without delay to preserve alveolar oxygenation in the event of difficult or failed TI (E.S. 85.7%).

2 G SADs have most of the ideal characteristics: easy insertion, high oropharyngeal sealing pressures, and they allow TI and gastric decompression.<sup>49,239,244</sup> They have thus shown superior performance to first generation devices and are more suitable for advanced use and as rescue devices.<sup>244-246</sup> Therefore, the immediate availability of an 2 G SAD is recommended, and to have the necessary competence to use it in all AW management sites (E.S. 100%).

The SAD to rescue a DAW should be selected prior to the procedure. Those with high first-attempt ventilation success and that allow IT through them<sup>238,247</sup> are of first choice: i-gel™ (Intersurgical Ltd., Wokingham, UK), Ambu® AuraGain™ (Ambu A/S, Ballerup, Denmark), LMA® Protector™ (The Laryngeal Mask Company Limited, Mahé, Seychelles), and iLTS-D (VBM Medizintechnik GmbH, Sulz, Germany).

One optimal attempt or a maximum of three attempts before declaring plan failure is recommended as the success rate decreases significantly with successive attempts<sup>248,249</sup> and these attempts increase trauma and delay transition between plans. During insertion of an SAD, cricoid pressure should be released if it is being used (E.S. 80%). Each attempt should include a change to improve the chances of success. Between and during attempts it is recommended to continue with peri-procedural oxygenation methods.

Rapid insertion and correct placement of the SAD are important to ensure the AW.<sup>250</sup> Rotation of 90°, jaw-thrust and the use of DL or VL (of choice) with the insert-detect-correct-as-you-got technique increase the efficacy and safety of the SAD by facilitating insertion, increase the first-attempt success rate and reduce pharyngeal trauma (E.S. 82.9%),<sup>246,250-257</sup> and can prevent malpositioning.<sup>257,258</sup> In contrast, blind insertion results in 50%-80% malpositioning<sup>246,256,257</sup> associated with suboptimal AW control, leakage or obstruction, increased risk of further displacement and increased morbidity.<sup>251,255,259</sup> Mal-positioned SADs are 26 times more likely to cause gastric insufflation and subsequent aspiration.<sup>255</sup> Table 4 shows the requirements for ideal positioning of an SAD, causes of malpositioning, and treatment options.<sup>246,255</sup> The recent videolaryngeal masks (VLM)<sup>259</sup> allow placement under direct view and immediate corrective manoeuvres without the aid of an additional device,<sup>66</sup> although there is still insufficient evidence.<sup>259-264</sup>

Correct placement of an SAD is clinically confirmed by a normal capnography waveform (ventilation grade 1) and maintenance of adequate alveolar oxygenation with peak inspiratory pressures of 20 cmH<sub>2</sub>O, oropharyngeal leak pressure >25 cmH<sub>2</sub>O, pulmonary auscultation, and passage of a tube through the gastric channel as complementary signs.<sup>259</sup>

Achieving effective ventilation and oxygenation provides time to stop and decide how to proceed according to the degree of urgency and type of procedure:

**Table 4** Requirements for ideal position of a supraglottic airway device, causes of malpositioning, and options for AW management.

**Correctly placed SAD (good seal and no leak)**

**Requirements for ideal position**

1. Tip of distal cuff in oesophagus
2. Epiglottis resting on the outside of the cuff
3. Epiglottic tip aligned with proximal cuff
4. Cuff adequately inflated to produce a seal (pressure 40–60 cm H<sub>2</sub>O)
5. No crease in the cuff (silicone is better than PVC)

**Avoid:**

- Hyperinflating the cuff (displacement of the SAD)
- Hyperinflating the cuff (risk of aspiration)
- SAD too deep/too small
- SAD too superficial/too large

**Malpositioned SAD confirmed with videolaryngoscopy<sup>255</sup> (leak and AW obstruction)**

**Causes**

1. Tip of distal cuff
  - a. Bending upwards
  - b. Bent backwards
  - c. Located between the vocal cords
2. Epiglottis in the bowl of the SAD
  - a. Without folding upwards
  - b. Folded downwards
  - c. Folded epiglottis

**Treatment options:**

- Jaw thrust to open the oropharyngeal space (increase the distance between the epiglottis and the posterior wall of the oropharynx) with the aim of repositioning the SAD<sup>268</sup>
- Different size or type of SAD (PVC cuff to silicone or reinforced tip)
- Guided technique with a bougie or orogastric tube<sup>269</sup>
- Magill forceps<sup>270</sup>

SAD: supraglottic airway device.

Adapted from Van Zundert AA, et al.<sup>246</sup>

- 1 If the situation is non-urgent (e.g. elective surgical procedure), the safest option is to awaken the patient and perform surgery under regional anaesthesia or postpone the intervention to perform an ATI.
- 2 If the situation is an emergency in (a) urgent surgery: it may be decided to continue the procedure while maintaining the second-generation SAD, but this is a high-risk option because the AW patency may be compromised in the course of the surgery; (b) the critically ill patient: a definitive AW is likely to be required, and therefore a FONA (tracheostomy) may be performed in anticipation of life-threatening hypoxaemia.
- 3 In intermediate cases, FOB-guided TI via the SAD may be chosen if the situation is stable, under adequate NMB and if the operator is competent (E.S. 97.1%, success rates close to 100%<sup>238</sup>). Blind TI is not recommended as the success rate is low (10%–20%), repeated attempts are required, and may result in increased trauma and impaired oxygenation.<sup>265,266</sup> The use of VLM allows TI without the help of FOB,<sup>66</sup> and many therefore shorten the technique and be especially advantageous in settings where FOB is not available, such as pre-hospital care and emergency care.<sup>267</sup>

Grade 2–3 ventilation or ineffective oxygenation after exhausting attempts requires immediate progression to a new plan.

### Face mask ventilation

FMV is an essential transitional technique during induction and treatment of an emergency AW and an indispensable rescue plan when other techniques fail.<sup>271</sup>

The presence of predictors of difficult FMV, and its use during emergency AW or as rescue from failed plans, make it especially advisable to use the optimal technique for FMV (triple manoeuvre of full neck extension, anterior jaw thrust, and mouth opening) from the outset, oropharyngeal or nasopharyngeal cannula placement,<sup>272</sup> and two-handed V-E technique,<sup>271,273</sup> either with two operators or with pressure-controlled ventilation with ventilator or other device,<sup>274,275</sup> in a patient with optimal positioning and intense NMB<sup>271,276–280</sup>) (E.S. 80%) in order to limit AW obstruction and optimise sealing to achieve grade 1 alveolar ventilation without causing gastric insufflation.<sup>126,275,281</sup> This also speeds up the transition between plans and reduces

peak ventilation pressure.<sup>282</sup> Face mask ventilation with modified triple manoeuvre is recommended over the E-C technique for the general population (1 C).

An effective attempt should be defined, in addition to a stable or improved normal SpO<sub>2</sub>, an acceptable tidal volume and AW pressure (4–5 mL/kg<sup>-1</sup> and <15–20 cmH<sub>2</sub>O, respectively<sup>273,275,283</sup>) and clinical criteria, by the presence of plateau phase on the capnography trace.<sup>284</sup> The use of scales to stratify the difficulty of FMV objectively, such as that proposed by the Japanese Society of Anaesthesiologists (Fig. 2, Part I), is recommended. This allows early declaration of failed FMV before desaturation (late sign) occurs.<sup>275</sup>

FMV is a dynamic procedure that requires continuous assessment until the AW is assured.<sup>285</sup> Declaration of failed FMV implies immediate transition to ventilation with SAD (E.S. 85.7%). Clinical deterioration and worsening oxygenation should prompt declaration of a CICO situation and immediate transition to FONA if ventilation with SAD has also failed.

A CICO situation should be declared using clear language that is easily understood by the entire team to create a shared model that facilitates an effective transition to FONA.<sup>119,286</sup>

## Recommendation

Face mask ventilation is recommended with modified triple manoeuvre over the C-E technique for the general population.

**Strong recommendation; low level of evidence**  
(⊕⊕⊖⊖)

## Front-of-neck access techniques

Failure of the three non-invasive supraglottic plans (primary and rescue) in the apnoeic patient requires declaration of a CICO situation and prompt FONA regardless of the SpO<sub>2</sub> value (E.S. 90.6%) because, in these circumstances, deterioration will be imminent if it is not already apparent. In the hospital setting, SEDAR SEMES SEORL-CCC recommends requesting the presence of an ENT specialist (or surgeon trained in performing tracheostomy) as soon as a CICO situation has been declared, although no procedure should be delayed until such time.

Delay in FONA is a major cause of morbidity and mortality.<sup>99,234,247,287,288</sup> Situational awareness and shared decision making, as well as good technical and HF training, can remove these psychological barriers to abandoning further attempts at non-invasive techniques.<sup>99,119,287,289–291</sup> The use of cognitive aids speeds up the transition to FONA by mitigating stress<sup>292</sup> (Fig. 1, Part I).

Psychological and technical preparation is recommended before declaring a CICO situation. A double set-up may be useful<sup>293</sup>: one team manages the AW conventionally while a second team is prepared and ready to perform a potential FONA. The multidisciplinary approach is associated with bet-

ter first-attempt success rates and faster time to securing the DAW.<sup>104</sup>

It is recommended to continue 100% supraglottic oxygen delivery, ensure intense NMB<sup>127,294</sup> and full neck extension for better laryngeal exposure.<sup>295</sup>

Emergency FONA encompasses three techniques: percutaneous cricothyrotomy, surgical cricothyrotomy, and surgical tracheostomy.

Cricothyrotomy is the technique of choice in a CICO situation (S.D. 91.4%) due to its relative simplicity, rapidity, high success rate, and low complication rate,<sup>100,296</sup> although the presence of an experienced ENT specialist or surgeon ensures that tracheostomy is performed as effectively and safely as a scalpel cricothyrotomy,<sup>297,298</sup> and therefore if the ENT specialist is present, they will lead the FONA using the technique they consider most appropriate.

The approach to a cricothyrotomy can be surgical or percutaneous. The evidence is limited as to which FONA approach is more appropriate in this setting. The few comparisons that have been made were in a simulation model and an animal model,<sup>99</sup> and therefore no one technique can be identified as superior to the others.<sup>99,101,299–302</sup> However, surgical access is successful in virtually 100% of cases.<sup>234,287,303</sup>

SEDAR SEMES SEORL-CCC recommend the scalpel-bougie-tube surgical cricothyrotomy method (E.S. 91.4%) for the following reasons<sup>98,234,247,287,291,294,301,304,305</sup>: (1) universality of the material, (2) it can be performed by a single operator, (3) technical competence is easy to acquire, (4) its execution, despite involving a greater psychological challenge,<sup>300,306,307</sup> it is characterised by its simplicity and speed, (5) it allows insertion into the trachea of an ETT or cannula with balloon and sufficient internal diameter to definitively secure the AW, protect it against aspiration, and allow effective gas exchange by means of PPV with conventional equipment and confirmation with capnography, and (6) its safety and efficacy profile. The required equipment includes a scalpel with blade no. 10, 20, or 21, ETT, or cannula with an internal diameter of no more than 6 mm and a bougie.

Other cricothyrotomy methods are considered valid, provided the appropriate expertise and equipment are available. Percutaneous cricothyrotomy approaches that achieve re-establishment of oxygenation with adequate ventilation are those that allow the introduction of a tube or cannula with an internal diameter  $\geq 4$  mm and 15 mm connector, or wide cannula techniques.<sup>247,291,294,301,305,308</sup> SEDAR SEMES SEORL-CCC leaves its use to the operator's discretion as first choice or as rescue due to failure of another method,<sup>309</sup> and therefore recommends acquiring skills in more than one technique.<sup>300,302,310</sup> Experience, availability of the equipment, and patient characteristics play a relevant role in the selection of technique.<sup>99,101,294,297,309,311</sup> The percutaneous technique, because it is more familiar and less intimidating, could be initiated earlier.<sup>300</sup>

It is essential to identify the CTM for the success of the technique. Detection by palpation has a high error rate. The use of ultrasonography over palpation is recommended to identify the cricothyroid membrane (1 C), and therefore SEDAR SEMES SEORL-CCC recommends acquiring the necessary skills. The laryngeal handshake technique, although slightly more time-consuming than conventional

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

techniques, has a higher success rate.<sup>312,313</sup> For cervical spine anatomies with anatomical landmarks that are difficult to identify by palpation, a vertical skin incision of more than 4 cm in the midline of the neck in a distal-proximal direction above the sternal notch is suggested to find the relevant anatomy.<sup>98,247,314</sup> Ultrasonography is superior to palpation in identifying the CTM and its immediate availability improves success.<sup>315–319</sup> However, it can be time-consuming, and therefore in a CICO situation it is only recommended if there is immediate availability and sufficient training.

After ensuring FONA, correct ventilation and alveolar oxygenation should be verified by capnography waveform, clinical assessment, pulse oximetry, and arterial blood gases when indicated.<sup>294</sup> A fibrobronchoscopy and radiological examination can complete the examination. Emergency cricothyrotomy should be converted to an ETT or tracheostomy as there is insufficient evidence as long-term treatment (E.S. 85.7%).<sup>296,320</sup> Conversion within 72 h is recommended to avoid the development of subglottic stenosis.<sup>301,321</sup>

Failure of a cricothyrotomy makes tracheostomy by a skilled operator advisable (E.S. 94.3%).<sup>100</sup>

CICO situations accounted for 39% of critical events and 25% of all anaesthesia-related deaths in the NAP4.<sup>287</sup> FONA is the last resort when non-invasive strategies have failed and is therefore of vital importance.<sup>322</sup> It is recommended that every institution standardise the technique and have a commercial or pre-assembled cricothyrotomy kit<sup>99,247,311,323</sup> with transparent packaging, which allows familiarisation with its contents and mental rehearsal of its use, and easy access, ideally located on the DAW trolley, available in every AW management site. All AW management professionals must acquire and maintain the necessary competence to perform a surgical or percutaneous cricothyrotomy with Seldinger's technique (E.S. 100%).<sup>99,101,119,247,291,298,301</sup> FONA should be feasible in any AW management site (E.S. 100%).<sup>324</sup>

## Recommendation

Ultrasonography is recommended over palpation to identify the cricothyroid membrane.

**Strong recommendation; low level of evidence**  
(⊕⊕⊖⊖)

## Pre-hospital setting

AW management in the pre-hospital setting is critical to patient survival and represents a real challenge for the emergency medical services (EMS).<sup>325,326</sup> In each scenario, different combinations of factors and constraints come together to create a high level of uncertainty that can turn any ordinary AW into a DAW.<sup>327</sup> Also, patient and operator positions may not always conform to those described in the manuals (subject trapped, confined, crushed, buried or in a place with simply no space). Survival depends on the cor-

rect interlocking of the entire chain of care. Table 5 shows the differential characteristics of AW management in the pre-hospital setting.<sup>328,329</sup>

## Human and ergonomic factors

A standardised rapid sequence induction (RSI) procedure is recommended to optimise outcomes, alleviate cognitive load in high-pressure situations, and improve technical and non-technical performance. Therefore, it is desirable to standardise the AW bag, have an ergonomic layout, and a checklist.<sup>330</sup> Fig. 4 shows the bag and ergonomic layout suggested by SEDAR SEMES SEORL-CCC for the pre-hospital setting.

## AW case/bag

Its objective is to store AW material in a standardised way, with a modular design, to keep it visible, accessible in less than a minute and organised in specific compartments for each of the four treatment plans, separated by Velcro® or zips, and marked with easily recognisable pictograms.<sup>331</sup> There should be at least one device for each plan (Fig. 4).

For outdoor work, it shall be a transportable, non-rigid case, bag, or rucksack, of reasonable size and weight, labelled, and manufactured in a different colour with approved material that is waterproof, resistant to corrosive agents, and easy to wash and sterilise. It shall have both handles and shoulder straps to free the hands and allow unimpeded access to the patient. Equipment should preferably be checked daily and at shift changes against a checklist.

## Ergonomics of positions during DAW management

Critical patient care in the pre-hospital environment is based on teamwork, the team usually consisting of a physician, a nurse, and one or more technicians. Therefore, there is usually only one operator.

The multiple scenarios, risks, patients, positions, and obstacles make it necessary to adapt to the setting. Each intervention is unique and unrepeatable. The situation requires accessing and controlling the scene and identifying the most serious subject, determining priorities, and establishing more than one approach plan. Most often the patient is found lying on the ground and a safety roll is possible, although not always. TI on the floor is a predictor of DAW.<sup>332</sup> Abnormal positions generate varying degrees of difficult-to-access DAW/AW, necessitating awareness of the anatomical changes and pathophysiology involved (Fig. 5). However, in an ideal scenario, SEDAR SEMES SEORL-CCC suggests the arrangement of team members as shown in Fig. 4. The physician controls ventilation and AW at the patient's head with a technician on their right to support AW management (Sellick, backward, upward, right lateral position[BURP], AW enlargement hook, traction support, supply of material, etc.). The nurse and second technician are positioned next to the individual's left shoulder/arm; while the AW case is opened to

**Table 5** Differential characteristics of AW management in the prehospital setting.

Body of doctrine	<ul style="list-style-type: none"> <li>Lack of a specific body of doctrine and a unified training model</li> </ul>
Operational procedures	<ul style="list-style-type: none"> <li>Research presents complex methodological challenges</li> <li>Highly complex clinical safety, quality control, and evaluation procedures</li> </ul>
Safety	<ul style="list-style-type: none"> <li>Limited level of evidence</li> <li>Protocols incomplete or with ostensible limitations (lack of cognitive aids and poor updating)</li> </ul>
Setting	<ul style="list-style-type: none"> <li>Usual setting of particular complexity (contextual DAW)</li> <li>Patient's home, public thoroughfare, or inside an ambulance</li> <li>Confined and/or inaccessible spaces</li> <li>Adverse context (infinite combinations): adverse weather conditions, excessive or poor lighting, noise level, social pressure</li> <li>Difficult access. Sometimes it will be necessary to manage the AW in abnormal positions</li> <li>High stress level</li> <li>Highly complex clinical context</li> </ul>
Patient and pathology	<ul style="list-style-type: none"> <li>Emergency indication: imminent AW compromise, absence of spontaneous ventilation, impaired level of consciousness, or agitation</li> <li>Physiological DAW: haemodynamic compromise, shunt, <math>V/Q</math> mismatch, ↓ FRC, full stomach</li> <li>Anatomical DAW</li> <li>Unknown history and clinical history</li> </ul>
Operator	<ul style="list-style-type: none"> <li>Single, less experienced operator due to multidisciplinary training</li> <li>Only one opportunity to control the AW (oedema, haemorrhage in AW...)</li> </ul>
Material and devices	<ul style="list-style-type: none"> <li>Devices not adapted to the pre-hospital setting (extreme temperatures, lighting)</li> <li>Lower solvency (e.g., portable aspirators with lower power)</li> <li>Limited availability</li> </ul>
Airway maintenance	<ul style="list-style-type: none"> <li>Medical transport carries added risks that require specific anticipation and planning: <ul style="list-style-type: none"> <li>- Accidental extubation in uncontrolled environment</li> <li>- Clinical and haemodynamic deterioration: early, delayed, and late</li> <li>To avoid complications, specific checklists are recommended for initiation of medical transport</li> <li>Contingency plans are required considering special circumstances (non-stop itineraries, air transport, limited space, or difficult access)</li> <li>Capnography is the guiding element</li> </ul> </li> </ul>

AW: airway; DAW: difficult airway; FRC: functional residual capacity;  $V/Q$  mismatch: ventilation/perfusion mismatch.

the right, and the monitor at the feet for proper visualisation. In case of cardiorespiratory arrest (CRA) the monitor should be placed on the subject's left shoulder for better handling.

For anticipated DAW, it is also recommended that teams have experience in awake TI, although the indications are narrower in emergency TI.<sup>336,337</sup>

## Pre-procedural assessment and planning

DAW is by definition not anticipated in the pre-hospital setting. However, pre-procedural assessment remains essential to anticipate potential difficulties and plan treatment.<sup>331</sup> Predictive tests can be difficult to perform, and, in many cases, there is no time for more than one second look. This rapid assessment can be useful but, where possible, should be combined with other tests.<sup>333</sup>

Different series have described as independent predictors of DAW: (1) AW obstruction, intubation, on the floor, and hyoid-mental distance less than three fingers,<sup>332</sup> (2) limited space at the scene, short neck, obesity, face and neck injuries, mouth opening less than 3 cm and ankylosing spondylitis,<sup>334</sup> and (3) Glasgow scale > 3, limited cervical spine movement, trismus/jaw tension, inability to palpate landmarks in the neck, presence of blood or vomit in the airway.<sup>335</sup>


## Peri-procedural oxygenation

Apnoeic oxygenation significantly reduces hypoxaemia during emergency TI,<sup>338,339</sup> making it, together with pre-oxygenation, essential for TI in the pre-hospital setting.<sup>340</sup> Standard nasal cannulae at 15 L/min should be included from the pre-induction period until TI in daily clinical practice except in patients with epistaxis, severe head trauma with possible skull base fracture, or complex facial fractures because it could worsen TI conditions and cause pneumocephalus.<sup>331</sup>

Any factors that may influence the success of TI should be optimised prior to the first attempt to achieve the best outcome.

## Rapid sequence induction

RSI is the most commonly used method in the pre-hospital setting. However, alternative techniques are of interest.<sup>335</sup> It is advisable to follow a standardised RSI procedure using



**Figure 4** Location of equipment and material in an ideal pre-hospital scenario.

N: nurse; P: physician, Mo: monitor; T: technician; DAW: difficult airway.

a checklist including drugs, calculation of dose and all available elements.<sup>341</sup>

### Cardiopulmonary resuscitation

Any abnormal situation increases the delay in ventilation and alveolar oxygenation, interrupts chest compressions, and delays recovery of spontaneous circulation (ROSC).<sup>342</sup>

The available evidence does not show strong differences between different non-invasive AW management plans.<sup>343-345</sup> Outcomes are conditioned by TI success rates. Therefore, if the desired level of TI efficacy is not achieved, ventilation and alveolar oxygenation should be given preference over the specific treatment plan, taking care not to interfere with other techniques (chest compressions, defibrillation, and treatment of potentially reversible causes of cardiac arrest).<sup>328</sup>

### Severe trauma

Severe head trauma carries a high risk of AW obstruction, pulmonary aspiration, hypoxia, brain injury and death.<sup>346</sup> Severe head trauma carries a high risk of AW obstruction, pulmonary aspiration, hypoxia, brain injury, and death. Pre-hospital TI is beneficial when performed by experienced physicians following standardised protocols.<sup>346,347</sup> Pre-hospital TI together with air transport could reduce overall mortality by 47%.<sup>326</sup>

Several factors may hinder AW management such as, (1) the possible presence of an unstable cervical spine injury necessitating bimanual alignment, (2) contaminated AW, flooded or pooled by tissue, vomitus, secretions, blood (bloody airway), etc., these conditions require aggressive management with manual release of fragments and radical aspiration of secretions using the suction assisted laryngoscopy airway decontamination (SALAD) manoeuvre,<sup>349</sup> (3) uncooperative or agitated patient, (4) fractures of



**Position 1. Operator kneeling.**

Allows good application of force by the left hand, although requires the operator to have dorsal flexibility. However, poor axis alignment may make laryngoscopy and oesophageal intubation difficult.



**Position 2. Operator in prone position.**

The axes are much more favourable. The left arm reduces its lever arm and, therefore, the axial traction force.



**Position 3. Operator seated at the patient's head.**

Good application of force. Poor axis alignment. Useful for narrow spaces.



**Position 4. Operator seated at the patient's side.**  
Very complex.



**Position 5. Operator straddling the patient's chest (reverse frontal).**

Requires change of hands. Allows good application of force. Axis alignment is not good. Usually requires support to maintain balance.




**Position 6. Operator in right lateral decubitus.**

Very complex. While the position reduces the power of the left arm, supporting the right arm reduces its level of precision. Useful for left-handers, with left-handed blades



**Position 7. Operator in left lateral decubitus.**  
**Paris SAMU position.**  
Very interesting. Increases the traction power of the left arm, which also has a lateral traction component due to the position, and further opens the airway. The right hand retains its precision.



**Position 8. Operator above and behind the patient.**  
Requires flexibility. Laryngoscopy is simple. Axis are good. Support is often required.  
With the patient upright, the need for gravity traction is reduced



**Position 9. Operator facing the patient.**  
Requires a change of hands. Enables application of force and laryngoscopy

**Figure 5** Main models of the approach to the difficult-to-access AW.

the facial mass, and (5) direct trauma to the AW, both penetrating and blunt (AW burn and/or inhalation syndrome). The presence of penetrating cervical trauma is the most frequent indication for awake TI while those

with maxillofacial trauma have the highest incidence of FONA.<sup>348,350</sup>

In these conditions, TI with RSI and VL with hyperangulated blade and preconfigured stylet is recommended.<sup>348,351</sup>

The impact of sunlight on the screen or the presence of blood or vomitus in the AW may condition the procedure.<sup>352</sup> If VL is not available, an SAD or DL with traction reduction can be used.<sup>329</sup> The use of DL may increase the risk of cervical spine injury.<sup>348,350</sup>

## Trapped, buried, crushed, or inaccessible patient

This is the paradigm of the difficult-to-access AW. This new field forces teams to expand their knowledge so as to take part in life support manoeuvres for rescue (8 methods) and in hostile environments.<sup>353</sup>

## Cuff pressure monitoring

Much laryngotracheal morbidity is related to inadequate cuff pressure.<sup>354</sup> Underinflation may cause hypoventilation or an increased risk of aspiration, while excessive pressure, even for short periods, may cause hoarseness, sore throat, impaired ciliary motility, and injuries such as mucosal swelling and ischaemia, laryngeal oedema, ulceration, stenosis, tracheoesophageal fistula, tracheomalacia, tracheal rupture, vocal cord paralysis, or nerve injury.<sup>241,355-357</sup> The incidence of these complications has reduced since the introduction of low-pressure, high-volume cuffs<sup>358</sup>; however, the devices used still cause avoidable harm.<sup>94,359</sup>

Intermittent cuff pressure monitoring by manometry is desirable after cuff pressure is established and periodically during maintenance<sup>360-364</sup> (not applicable in crisis situations). The use of continuous monitoring devices to constantly maintain the cuff pressure in range in the critically ill patient may decrease the risk of ventilator-associated pneumonia.<sup>365-367</sup> Continuous monitoring with cuff pressure manometry is suggested (1 C).

Cuff pressure should be set at the minimum pressure necessary to ensure an effective and safe seal. The pressure should remain between 20 and 30 cmH<sub>2</sub>O (ideally less than 25 cmH<sub>2</sub>O) for ETT and tracheostomy and cricothyrotomy tubes, and < 60 cmH<sub>2</sub>O for SADs (E.S. 94,3%),<sup>241,354,358,368-371</sup> from insertion to removal. If within these limits the seal is inadequate, it may be necessary to reposition the device or resize it.<sup>372</sup>

Nitrous oxide diffuses into the cuff,<sup>359</sup> requiring further measurements after 20 min, when the pressure stabilises, and again if its concentration increases.<sup>373</sup>

## Recommendation

Continuous cuff pressure monitoring with manometry is suggested.

**Strong recommendation; low level of evidence**  
(⊕⊕⊖⊖)

## Extubation

Tracheal extubation is a high-risk procedure with major physiological implications.<sup>374</sup> It may evoke a haemodynamic response to stress, cough, laryngospasm, or agitation and, secondarily, an increase in intracranial or intraocular pressure.<sup>375-378</sup> Protective laryngotracheal reflexes remain impaired for several hours after extubation, facilitating aspiration. Failure of extubation occurs when a patient is unable to maintain oxygenation, adequate ventilation, clearance of secretions, or AW patency and can have catastrophic consequences, particularly in those with DAW.<sup>376</sup> AW obstruction is its main cause and is often associated with post-obstructive pulmonary oedema with severe hypoxia.<sup>378</sup> Failed extubation, generally defined as the need for reintubation within 24–72 h, occurs in approximately .1%–.45% of general anaesthesia<sup>379</sup> and is 10 times more prevalent in patients with sleep apnoea-hypopnoea syndrome (SAHS), or those undergoing AW procedures, and its prevalence is 10-fold higher in critical care areas.<sup>380</sup> This is an event with potentially severe outcomes.<sup>381-383</sup> Thus, secondary cases of death or permanent brain damage constitute one third of all anaesthesia-related cases.<sup>288,384</sup> Poor anticipation and planning for ät-riskextubation are key issues.<sup>288,384</sup> Extubation is an elective procedure that must be prepared for, followed by a stepwise strategy and meticulous follow-up.<sup>375,376,378,385</sup> Fig. 6 shows a cognitive aid for planning extubation based on risk.


The primary goal in tracheal extubation is to preserve alveolar oxygenation in addition to avoiding reintubation. Any reintubation can be considered potentially difficult as it involves additional complexity (E.S. 97.1%, distorted anatomy with restricted access, secretions, blood or oedema, limited time, and in a high-stress environment).<sup>375,376,385</sup> Risk stratification based at anatomical, physiological, and contextual level<sup>376,383,386,387</sup> can determine the chances of success conditioned by extubation tolerance and reintubation feasibility, establishing an individualised strategy and optimising AW-related factors and physiological status such as hypoxaemia, hypercapnia, residual NMB, hypothermia, or oedema in the AW.<sup>355,383,387</sup> Leak testing,<sup>388</sup> preferably quantitative (leak volumes <110 mL or <12%–24% of tidal volume determine reduced AW patency and risk of post-extubation stridor due to laryngeal oedema<sup>387,389</sup>), ultrasound assessment,<sup>390,391</sup> and visualisation of the larynx with VL or FOB can help in decision making (E.S. 97.1%) because they enable assessment of AW patency and determining the presence of periglottic oedema or bleeding prior to extubation,<sup>355</sup> although they are not a specific predictive test for successful extubation.<sup>392</sup>

Once the decision has been made to proceed with extubation, the first step is to review the original TI and update the AW assessment, and general risk factors.<sup>383</sup> Careful preparation beforehand, conveying potential problems and early warnings, and establishing appropriate rescue plans for oxygenation and reintubation in case of failure of the primary plan are conducive to safe and successful extubation.<sup>378,383</sup> Important prior to intervention<sup>378</sup> are (1) pre-oxygenation until extubation, (2) oropharyngeal suctioning of secretions or blood under direct visualisation, ideally with laryn-



# Risk-based planning for tracheal extubation

## Assessment of risk factors




## Planning

### Single category RF

- No RF. → Awake vs. asleep
- Pre-existing anatomical RFs
  - Patent AW → Awake + EC
  - AW obstruction → Assess tracheostomy
- Physiological RFs → Pharmacological adjuvants  
Bailey manoeuvre
- Acquired RFs
  - Reversible → Defer
  - Late resolution or irreversible → Assess tracheostomy
- Contextual RFs → Defer

### Combine RFs



**Figure 6** Cognitive aid for planning, risk stratification, and decision making for AW extubation.

Adjts: adjutants; AW: airway; DAW: difficult airway; EC: exchange catheter; RF: risk factors; SAHS: sleep apnoea-hypopnea syndrome; TI: tracheal intubation.

goscopy, to avoid soft tissue trauma, (3) placement of a bite block,<sup>386</sup> (4) correct positioning of the patient, (5) reversal of NMB, neuromuscular monitoring in combination with reversal agents aiming for a train-of-four ratio  $\geq .90$  is essential to avoid residual NMB,<sup>393-396</sup> (6) minimisation of head and neck movements and reduction of noxious stimuli to avoid laryngospasm, (7) emergence to state of wakefulness,<sup>377,378,385-387</sup> deep extubation is an inappropriate technique for DAW or when there is risk of aspiration,<sup>377,386</sup> (8) apply positive pressure, deflate the cuff and remove the ETT, (9) administer 100% oxygen, confirm AW patency and adequate spontaneous ventilation, and (10) continuous oxygen therapy until full recovery, surveillance, monitoring, and skilled assistance to cope with potential emergency tracheal reintubation.<sup>385</sup> The DAW team must be prepared and remain immediately accessible.

Prophylactic administration of corticosteroids prior to elective extubation is associated with a significant decrease in the incidence of post-extubation adverse events in the AW and reintubation, therefore patients at high risk of obstruction may benefit.<sup>397-399</sup> In those with absent or reduced leak, administration of a corticosteroid at least four hours prior to extubation is recommended.<sup>376,387,389</sup> Prophylactic administration of corticosteroids prior to extubation is suggested in patients at high risk of obstruction (1 B).

There are several advanced methods for at-risk extubation that should only be performed by experienced personnel.<sup>378</sup> The administration of pharmacological adjuvants<sup>400-405</sup> such as a remifentanil infusion<sup>406</sup> or the Bailey manoeuvre, which involves placing a SAD over the ETT and then removing it,<sup>407</sup> may be considered when gentle emergence and attenuation of undesirable cardiovascular or respiratory responses is required.<sup>408-410</sup> The use of VLMs for TI may have a role to play in safer and simplified TI because they allow the switch to be performed by simply pulling back the ETT through its ventilation channel, vision-guided, and reversing it instantaneously.<sup>66</sup> Both techniques require a deep level to perform, and may therefore be inappropriate in patients in whom reintubation may be difficult or there is a risk of aspiration.<sup>1,378</sup>

Awake extubation using advanced techniques is the most appropriate method for DAW (E.S. 94.3%), because it maintains upper AW patency with re-established muscle tone, protective reflexes, and spontaneous ventilation.<sup>410</sup> Airway exchange catheters are the most commonly used adjuvant to salvage extubation in potential or known DAW.<sup>1</sup> These devices allow tracheal reintubation by acting as a guide over which the ETT can be reinserted under direct view<sup>185</sup> and insufflation of oxygen at low flows or jet ventilation through them in potentially life-threatening hypoxaemia, although this should be avoided as even low flows have caused barotrauma.<sup>411</sup> They should be maintained until reintubation is unlikely.<sup>376</sup> Their use has been associated with high overall (92%) and first-attempt (87 vs. 14%) success rates, fewer oesophageal intubations, and fewer episodes of desaturation, bradycardia, or hypotension.<sup>412</sup> The main risks of the technique are AW stimulation, trauma with subcarinal insertions, and accidental catheter dislodgement. In adults, they should never be inserted deeper than 25 cm from the lips.<sup>378</sup> Staged extubation kits consisting of guidewire and reintubation catheter<sup>413,414</sup> are associated with an overall success rate of 93%,<sup>415</sup> and appear to improve tolerance at

the cost of increased accidental dislodgement.<sup>416</sup> If reintubation over an intubation catheter is necessary, limiting the difference between the internal diameter of the ETT and the external diameter of the catheter, establishing an appropriate NMB, viewing advancement of the ETT by VL facilitate the procedure by avoiding impingement of the epiglottis and arytenoids.<sup>383,417</sup> The remaining recommendations for facilitating FOB-guided TI are equally applicable to exchange catheter-guided TI. Failure of the technique means referring to the unanticipated DAW algorithm.

Capnography should be available in recovery units and used in high-risk cases (E.D. 97.1%). Oxygen therapy with nasal goggles and masks with capnography line after extubation facilitates early detection of respiratory depression, hypoventilation, hypercapnia, or post-extubation AW obstruction.<sup>379,387</sup>

Strategies to prevent extubation failure include head tilt and supplemental oxygen administration. A good oxygenation strategy with NIV or HFNO may prevent reintubation in at-risk populations.<sup>374,418-421</sup>

## Recommendation

Prophylactic administration of corticosteroids prior to extubation is recommended in patients at high risk of AW obstruction.

**Strong recommendation; moderate level of evidence (⊕⊕⊖⊖)**

## Documentation

Documentation of AW management is necessary to guide future management. In addition to the collection of consent and pre-procedural assessment it should include methods of peri-procedural oxygenation, topicalisation, awake sedation, induction, device, adjuvants and ETT used, approach (right nasal, left nasal, or oral), number of attempts, extubation, and any difficulties or complications. Appendix A, supplementary material 5 shows a template for recording relevant AW-related information.

A history of failure of previous procedures is the most accurate predictor of failure of subsequent treatments (E.S. 97.1%).<sup>422</sup> Documenting the difficulty is one of the most important actions to prevent complications as it facilitates decision-making and allows for a targeted structured approach and efficient transition with less instrumentation by avoiding persevering with plans that have failed in previous procedures.<sup>422,423</sup> Therefore, it is appropriate to diversify the means of documenting a DAW as this increases the chances of successful transmission of this critical information.<sup>423,424</sup> Thus, the following are recommended<sup>11,294,424</sup>: (1) recording in the electronic history with detailed description and visual record, images and/or video, of the anatomy and the successful technique employed,<sup>425</sup> (2) verbal communication to the conscious patient and the family member, guardian, or caregiver, (3) written report to the patient, Appendix A, supplementary

material 6 includes the DAW notification template proposed by SEDAR SEMES SEORL-CCC, (4) notification bracelet or necklace or ID card with QR code for access to clinical information, Appendix A, supplementary material 7 shows the means of identifying a DAW suggested by SEDAR SEMES SEORL-CCC, (5) alert each time the electronic patient record is accessed, and (6) entry in national or international databases.<sup>426</sup> It is recommended to standardise records with mandatory fields and free text sections to include the characteristics of the AW, the nature of the difficulty, failed and successful techniques.<sup>294,424</sup> Structured notes in the electronic record have shown an improvement with reduction of adverse events.<sup>427,428</sup>

### Management in the field of the airway

In addition to individual commitment, a proactive state, and institutional and departmental involvement are required to implement strategies.<sup>429,430</sup> The action plan to improve safety, cost-effectiveness, and quality of care in AW management includes (1) standardisation of practice and equipment in all locations, effective adherence to guidelines, a standardised DAW trolley,<sup>431</sup> removal of barriers and application of facilitators,<sup>432-434</sup> (2) coordination of departments that manage the AW (anaesthetics, intensive care, ED, ENT, and pre-hospital care), (3) procurement and availability of appropriate equipment, (4) continuous staff training and education, (5) regular audits and error limitation programme: (a) incident analysis, (b) identifying and addressing departmental and institutional factors that may have contributed to the event for learning and implementation of improvements, (c) case review and discussion of alternative plans, (6) incident documentation and notification plans,<sup>435</sup> (7) DAW coding, recording, identification and alerting systems, and (8) organisational resource guidelines and continuous process improvement.<sup>436,437</sup>

Appointing an AW lead in each institution is recommended (E.S. 100%),<sup>11,429,430,438</sup> a professional with experience in the field, who liaises with management to assume the actions and organisational aspects indicated. Their primary objective is to provide each professional with the necessary tools.<sup>429,439</sup> Some hospitals have set up multidisciplinary teams of specialised staff with defined roles to increase safety.<sup>440-442</sup> The creation of a national network of AW leads and teams can enable the field to develop.<sup>438,439</sup>

Decisions on the procurement of AW devices should be supported by formal evaluation based on scientific evidence.<sup>24,443,444</sup>

### Teaching and training

AW management is an essential competence in anaesthesia, critical care, pre-hospital, and emergency medicine.<sup>445,446</sup> Poor training is a common causal or contributing factor to complications<sup>234,384</sup> and impacts on the confidence operators have in the use of devices

and performing essential techniques such as FOB and cricothyrotomy.<sup>447-449</sup> Optimising teaching and training is key to increasing safety.<sup>322</sup>

There is limited evidence and lack of standardisation in AW management teaching,<sup>324,450</sup> and therefore strategies are extrapolated from other fields. Good training should include cognitive skills based on theoretical principles and technical and non-technical skills.<sup>322,451</sup> Competency acquisition should be gradual, through a cognitive phase, simulation, and clinical training with problem solving until the learning curve is complete, with assessment and feedback from the instructor at each phase (E.S. 100%).<sup>437,452,453</sup> Competency-based education requires individualised instruction.<sup>454</sup> Training should follow a learning approach consisting of reaching a predefined standard goal of mastery of a skill to escalating levels of difficulty and stress.<sup>455</sup> It is a learner-centred, evidence-based method,<sup>456</sup> potentially associated with better outcomes.<sup>457,458</sup>

Education and training should cover all sections of the guidelines in a structured way,<sup>37,322,445,455</sup> with special priority given to core, versatile, and time-critical skills.<sup>445</sup> Standardisation of a teaching and training programme is highly recommended. Simulation plays a relevant role, mainly in the acquisition of non-technical skills such as teamwork, the incorporation of guidelines and cognitive aids in clinical practice, and the rehearsal of tasks and resolution of rare life-threatening events.<sup>437,448,455,459-464</sup> Guidelines and algorithms are training tools for gaining knowledge and learning strategies and act as a resource during debriefing,<sup>322,465,466</sup> while cognitive aids are a good implementation tool for rehearsal during simulation.<sup>127,322</sup> Structured debriefing improves clinical knowledge, skill acquisition, and implementation of skills in practice.<sup>455,467</sup>

Operators with greater professional experience may be at risk of providing lower quality care,<sup>446,468</sup> and therefore require ongoing education and regular training to develop the new skills needed with the addition of new devices or techniques, and to maintain competencies,<sup>322,445,455,469,470</sup> preferably on an annual basis (E.S. 97.1%).<sup>471</sup> A proactive organisational and individual attitude is necessary. It is highly recommended that each service designate a local AW lead to promote quality interprofessional and multidisciplinary training programmes, with defined objectives, evaluation and supervision, and to ensure dissemination and adherence to the guidelines and cognitive aids.<sup>322,324,439,445,448,472</sup> Training should be diversified to provide the operator with alternatives.<sup>43</sup> Appendix A, supplementary material 8 shows the skills recommended by SEDAR SEMES SEORL-CCC that should be included in any AW training programme and the training methods needed to achieve objectives.

### Summary of recommendations from the systematic literature search

Search strategies and GRADE tables are shown in the supplementary material.

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

No.	Recommendation	Level of evidence	Grade of recommendation
<b>Pre-procedural assessment and planning</b>			
1.	A diagnosis of SAHS is a predictor of difficult face mask ventilation	Low	Strong
2.	A diagnosis of SAHS is a predictor of difficult tracheal intubation	Moderate	Strong
3.	Gastric ultrasound examination is recommended to assess the risk of aspiration in at-risk situations	Low	Strong
<b>Preparation</b>			
4.	Capnography waveform is recommended as gold standard to confirm alveolar ventilation	Moderate	Strong
5.	Ramped position or elevated head to 30° in the obese population is recommended to improve tracheal intubation conditions	Low	Strong
6.	Ramped position prolongs safe apnoea time in obese populations	Moderate	Strong
<b>Peri-procedural oxygenation</b>			
7.	HFNO is recommended as first-line pre-oxygenation technique for patients with mild hypoxaemia	Low	Strong
8.	NIV is recommended over conventional oxygen therapy for anaesthetic induction in the obese patient	Moderate	Strong
9.	Oxygenation during apnoea with high-flow nasal goggles (NOT DESAT/HFNO) is recommended	Low	Strong
<b>Rapid sequence induction</b>			
10.	Neuromuscular block is recommended to improve TI conditions and the incidence of AW-related adverse events in the general population	Moderate	Strong
11.	The rocuronium + sugammadex combination is not inferior to succinylcholine for RSI	Moderate	Strong
<b>Unanticipated difficult airway</b>			
Tracheal intubation			
12.	The routine use of VL is recommended over DL as the primary device for TI	Moderate	Strong
13.	The use of a dynamic or articulated bougie (flex-tip or FOB type) versus a conventional stylet is recommended for TI in patients with difficult airway	Low	Strong
14.	Parker Flex ETT is recommended over conventional ETT for FOB-guided TI in the general population	Moderate	Strong
15.	Parker Flex ETT over conventional ETT for TI guided by FOB and laryngoscopy in the general population is suggested to reduce complications	Low	Strong
Face mask ventilation			
16.	Face mask ventilation is recommended with modified triple manoeuvre over the C-E technique for the general population	Low	Strong
Front-of-neck access			
17.	Use of ultrasonography over palpation to identify the cricothyroid membrane is recommended	Low	Strong
<b>Pressure cuff monitoring</b>			
18.	Continuous pressure cuff monitoring with manometry is suggested	Low	Strong
<b>Extubation</b>			
19.	Prophylactic administration of corticosteroids prior to extubation in recommended in patients at high risk of AW obstruction	Moderate	Strong

AW: airway; DL: direct laryngoscopy; ETT: endotracheal tube; FOB: fibrobronchoscopy; HFNO: high-flow nasal oxygen; NIV: non-invasive ventilation; NO DESAT: nasal oxygen therapy during efforts to secure an ETT; RSI: rapid sequence induction; SAHS: sleep apnoea-hypopnoea syndrome; TI: tracheal intubation; VL: videolaryngoscopy.

## Expert statement from the results of the Delphi questionnaire

No.	Question	% in favour [in favour; neutral; against]
<b>Human factors</b>		
1.	The number of attempts for each non-invasive treatment plan should be limited to three	88,6 [31; 2; 2]
2.	The first attempt should be made under optimal conditions	100 [35; 0; 0]
3.	The most appropriate primary technique should be the one that offers the best guarantee of first-attempt success	94,3 [33; 1; 1]
4.	Visual cognitive aids are recommended for the management of emergency crises	97,1 [34; 1; 0]
5.	A standardised difficult airway trolley is recommended in areas with AW management	100 [35; 0; 0]
6.	Use of checklists is recommended to reduce the incidence of human error, improve task completion time, and reinforce a culture of safety in the AW management	100 [35; 0; 0]
7.	The use of ergonomic and communication models is recommended	91,4 [32; 3; 0]
<b>Pre-procedural assessment and planning</b>		
8.	Pre-procedural assessment is recommended for all patients requiring AW management	100 [35; 0; 0]
9.	Pre-procedural assessment of AW should be multifactorial, structured and aimed at detecting anatomical, physiological, and contextual DAW	97,1 [34; 1; 0]
10.	AW exploration can start by detecting predictors of difficulty or failure for the primary plan and subsequently for the 3 alternative plans	97,1 [34; 1; 0]
11.	Multivariate models may have a higher predictive power	97,1 [34; 1; 0]
12.	Decision-making should be individualised according to patient, operator, context, and time	97,1 [34; 1; 0]
13.	Restriction of food and fluid intake following preoperative fasting guidelines is necessary	97,1 [34; 1; 0]
14.	The presence of a full stomach indicates that the AW should be protected with TI	88,6 [31; 2; 2]
<b>Preparation</b>		
15.	Capnography waveform should be available in all AW management sites to assess the success of any of the 4 plans used	97,1 [34; 1; 0]
<b>Basic options for difficult airway management</b>		
16.	Before each procedure, the appropriateness of the treatment should be assessed, and a risk-benefit analysis performed	97,1 [34; 1; 0]
17.	Awake management is recommended when there is a high degree of difficulty or impossibility of TI, combined predictors of difficulty, or physiological disturbances, and negative contextual conditions	82,9 [29; 5; 1]
18.	Induction of general anaesthesia with preservation of spontaneous ventilation is suggested in situations where awake management is recommended but general anaesthesia is unavoidable due to uncooperativeness or urgency, and there are no physiological or contextual predictors of difficulty or obstructive pathology	91,4 [32; 2; 1]
19.	When physiological or contextual predictors of difficult AW exist, the benefit of deferral may be assessed if it outweighs the risk of proceeding to airway management, or alternative anaesthetic strategies may be considered	85,7 [30; 5; 0]
<b>Known or anticipated difficult airway</b>		
20.	Awake management is the option of choice to ensure a known or anticipated DAW	85,7 [30; 4; 1]
21.	High-flow nasal oxygen therapy is recommended over conventional low-flow cannulae	91,4 [32; 3; 0]
22.	NIV with endoscopic mask may have a role in TI in the critically ill patient with hypoxaemia	82,9 [29; 6; 0]
23.	Premedication with an antisialogogue is recommended to optimise the efficacy of local anaesthetic efficacy and field of view, glycopyrrrolate being the preferred option	80 [28; 5; 2]

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

24.	Sedation is an optional adjunct to adequate topical anaesthesia in awake airway management	88,6 [31; 2; 2]
25.	The goals of conscious sedation for awake AW management are effective amnesia, patient satisfaction, and analgesia to reduce cough, gag, and haemodynamic reflexes, while preserving AW patency, spontaneous ventilation, and protective laryngeal reflexes	94,3 [33; 2; 0]
26.	If the selected primary technique (FOB or VL) fails, the alternative technique should be used	80 [28; 6; 1]
27.	A third attempt may benefit from a multimodal approach (VL + FOB)	100 [35; 0; 0]
28.	The combination of a TI SAD and FOB may be useful as a rescue technique to maintain oxygenation, maintain AW patency and to perform a TI through it	100 [35; 0; 0]
29.	A smaller than usual ETT is recommended with VL and FOB	85,7 [30; 4; 1]
30.	Decreasing the difference between the outer diameter of the FOB and the inner diameter of the ETT is recommended to facilitate FOB-guided TI	85,7 [30; 3; 2]
31.	Standard PVC ETTs are not recommended for FOB-guided TI as they are more likely to impact glottic structures	71,9 [23; 4; 5]
32.	After visual confirmation of TI, it is recommended to induce general anaesthesia after establishing cuff pressure and capnographic confirmation of TI	94,3 [33; 2; 0]
33.	Alternative techniques and approaches should be planned in advance and implemented without delay after failure of the primary approaches	100 [35; 0; 0]
34.	With a high probability of awake TI failure, it is recommended to prepare a FONA in parallel to the invasive treatment plan in case of total obstruction	88,6 [31; 4; 0]
35.	Awake tracheostomy under local anaesthesia is recommended in the presence of pre-existing critical AW compromise	82,9 [29; 6; 0]
36.	Awake cricothyrotomy would be the most indicated technique in the presence of emergent critical compromise	91,4 [32; 3; 0]
37.	Awake ECMO under local anaesthesia may be the safest option when all 4 conventional plans are expected to be impossible, unsuccessful, or ineffective, with risk of full AW obstruction	90,6 [29; 1; 2]
<b><i>Unanticipated difficult airway</i></b>		
Peri-procedural oxygenation		
38.	HFNO should be considered as the first-line pre-oxygenation technique for patients with mild hypoxaemia ( $\text{PaO}_2/\text{FiO}_2 > 200 \text{ mmHg}$ ), while NIV is the technique of choice in those with severe hypoxaemia ( $\text{PaO}_2/\text{FiO}_2 \leq 200 \text{ mmHg}$ )	87,5 [28; 3; 1]
39.	Pre-oxygenation with NIV + HFNO and apnoeic oxygenation with HFNO should be a priority option for critically ill patients during TI	85,7 [30; 4; 1]
Rapid sequence induction		
40.	RSI is the recommended technique when there is a significant risk of aspiration in an AW without predictors of difficulty	97,1 [34; 1; 0]
41.	RSI with or without Sellick manoeuvre is recommended for all emergency TIs	84,4 [27; 1; 4]
42.	The use of checklists is suggested for safe preparation of RSI	97,1 [34; 1; 0]
43.	Premedication with non-particulate antacid immediately before induction and an H <sub>2</sub> -receptor antagonist 40–60 min before, or a proton pump inhibitor to increase pH and reduce the volume of gastric contents is suggested in patients at high risk of aspiration	82,9 [29; 5; 1]
44.	Nasogastric tube treatment should be individualised	88,6 [31; 4; 0]
45.	Highly efficient suction devices with large multi-hole suction tubes should be available in case of potential regurgitation	100 [35; 0; 0]
46.	An elevated head position to 20°–30° is recommended to prevent passive regurgitation and, if regurgitation occurs, the Trendelenburg position, turning the head to one side and suctioning the oropharynx and trachea before initiating positive pressure ventilation	94,3 [33; 2; 0]
47.	The choice of hypnotic as well as the dose and speed of administration should be individualised	91,4 [32; 3; 0]
48.	Delayed sequence induction is suggested in agitated and uncooperative patients for adequate pre-oxygenation	71,9 [23; 3; 6]
49.	Routine use of cricothyroid pressure cannot be recommended	81,3 [26; 2; 4]

50.	A modified RSI can be applied in patients at high risk of hypoxia who are not candidates for ATI	85,7 [30; 5; 0]
<b>Tracheal intubation</b>		
51.	Devices with a standard Macintosh-type blade (allows direct and indirect laryngoscopy) are appropriate for AW management without predictors of difficulty, while those with a hyperangulated blade (without or with a guide channel) are indicated for known or anticipated DAW	94,3 [33; 1; 1]
52.	It is recommended that a bougie be available in all AW management sites	97,1 [34; 1; 0]
53.	Absence of capnography trace (ventilation grade 3) indicates failed TI until proven otherwise	80 [28; 6; 1]
54.	Capnography waveform monitoring during maintenance of mechanical ventilation is highly recommended in all sites	100 [35; 0; 0]
<b>Ventilation with supraglottic airway device</b>		
55.	A SAD should be inserted without delay to preserve alveolar oxygenation in the event of difficult or failed TI	85,7 [30; 3; 2]
56.	Immediate availability of a second-generation SAD is recommended, as well as competency in its use in all AW management sites	100 [35; 0; 0]
57.	Cricoid pressure should be released during insertion of an SAD if it is being used	80 [28; 5; 2]
58.	90° rotation, jaw-thrust, and DL or VL (of choice) with the insert-detect-correct-as-you-gotchnique increase the efficacy and safety of the SAD by facilitating insertion, increase first-attempt success rate, and reduce pharyngeal trauma	82,9 [29; 4; 2]
59.	FOB-guided TI through the SAD may be chosen if the situation is stable, under adequate NMB and if the operator is competent in the technique	97,1 [34; 1; 0]
<b>Face mask ventilation</b>		
60.	For FMV it is recommended at the beginning to use the optimal technique (triple manoeuvre of full neck extension, anterior jaw thrust, and mouth opening, placement of oropharyngeal or nasopharyngeal cannula and two-handed V-E technique, in a patient with optimal positioning and strong NMB	80 [28; 3; 4]
61.	Declaration of failed FMV implies immediate transition to VSAD	85,7 [30; 2; 3]
<b>Front-of-neck access</b>		
62.	Failure of all 3 non-invasive plans (primary and rescue), regardless of SpO <sub>2</sub> value, requires verbalisation of the need for and subsequent FONA	90,6 [29; 0; 3]
63.	Cricothyrotomy is the technique of choice in a CICO situation	91,4 [32; 2; 1]
64.	The scalpel-bougie-tube surgical technique is recommended for cricothyrotomy	91,4 [32; 2; 1]
65.	A FONA should be feasible wherever AW is managed	100 [35; 0; 0]
66.	Emergency cricothyrotomy should be converted to ETT or tracheostomy because there is insufficient evidence for it as long-term management	85,7 [30; 3; 2]
67.	Failure of a cricothyrotomy to secure the AW makes it advisable for tracheostomy to be performed by a skilled operator	94,3 [33; 1; 1]
68.	Any AW management practitioner must acquire and maintain the necessary competency to perform a surgical or percutaneous cricothyrotomy with Seldinger's technique	100 [35; 0; 0]
<b>Cuff pressure monitoring</b>		
69.	Cuff pressure should be established at the minimum pressure necessary to ensure an effective and safe seal. Pressure should remain between 20 and 30 cm H <sub>2</sub> O for ETT and tracheostomy and cricothyrotomy tubes, and < 60 cm H <sub>2</sub> O for SADs	94,3 [33; 1; 1]
<b>Extubation</b>		
70.	Any reintubation can be considered potentially difficult as its management involves additional complexity	97,1 [34; 1; 0]
71.	Leak testing, preferably quantitative, ultrasound assessment, and laryngeal visualisation with VL or FOB may facilitate decision making	97,1 [34; 1; 0]
72.	Awake extubation with use of advanced techniques is the most appropriate method for DAW	94,3 [33; 2; 0]
73.	Capnography should be available in recovery units and used in high-risk cases	97,1 [34; 1; 0]

**Documentation**

74.	A history of failure of previous procedures is the most accurate predictor of failure of subsequent management	97,1 [34; 1; 0]
Management in the field of the airway		
75.	Allocation of an AW lead in each institution is recommended	100 [35; 0; 0]
Teaching and training		
76.	Competency acquisition should be gradual, through a cognitive phase, simulation, and clinical training with problem-solving until the learning curve is complete, with assessment and feedback from the instructor at each phase	100 [35; 0; 0]
77.	Ongoing teaching and regular training are required for the development of new skills or techniques and the maintenance of competencies, preferably on an annual basis	97,1 [34; 1; 0]

AW: airway; CICO: can't-intubate-can't oxygenate situation; DAW: difficult airway; DL: direct laryngoscopy; ECMO: extracorporeal membrane oxygenation; ETT: endotracheal tube; FOB: fibrobronchoscopy; FiO<sub>2</sub>: fraction of inspired oxygen; FONA: Front-of-neck access; HFNO: high-flow nasal oxygen therapy; FMV: face mask ventilation; NIV: non-invasive ventilation; NMB: neuromuscular block; PaO<sub>2</sub>: partial arterial pressure of oxygen; PVC: polyvinyl chloride; RSI: rapid sequence induction; SAD: supraglottic airway device; 2 GSAD: second generation supraglottic airway device; SpO<sub>2</sub>: peripheral oxygen saturation; TI: tracheal intubation; VSAD: Ventilation with supraglottic airway device; VL: videolaryngoscopy.

**Authors' contribution**

- Manuel Á. Gómez-Ríos: drafting of the manuscript, preparation of all cognitive aids and graphic material, tables and annexes, literature review, critical reading, levels of evidence, final revision of the document.
- José Alfonso Sastre: Draft of RSI sections, SAD, and checklist, risk factor tables, information document models, literature review, critical reading, levels of evidence, final revision of the document.
- Xavier Onrubia-Fuertes: FONA contribution, unanticipated difficult tracheal intubation, literature review, final revision of the document.
- Teresa López: Draft of SAD and ECMO sections, literature review, critical reading, levels of evidence, final revision of the document.
- Alfredo Abad-Gurumeta: literature review, critical reading, levels of evidence, final revision of the document.
- Rubén Casans-Francés: literature review, critical reading, final revision of the document.
- David Gómez-Ríos: literature review, critical reading, final revision of the document.
- José Carlos Garzón: literature review, critical reading, final revision of the document.
- Vicente Martínez-Pons: Algorithm for unanticipated difficult tracheal intubation, literature review, final revision of the document.
- Marta Casalderrey-Rivas: literature review, critical reading, final revision of the document.
- Miguel Ángel Fernández-Vaquero: Algorithm for unanticipated difficult tracheal intubation, literature review, and critical reading aimed at AW predictors and assessment, final revision of the document.
- Eugenio Martínez-Hurtado: Algorithm for unanticipated difficult tracheal intubation, final revision of the document.
- Ricardo Martín-Larrauri: Algorithm for unanticipated difficult tracheal intubation, final revision of the document.

- Laura Reviriego-Agudo: Algorithm for unanticipated difficult tracheal intubation, literature review, final revision of the document.
- Uxía Gutierrez-Couto: literature search strategies.
- Javier García-Fernández: critical reading, final revision of the document.
- Alfredo Serrano Moraza: prehospital setting section, literature review, critical reading, final revision of the document.
- Luis Jesús Rodríguez Martín: prehospital setting section, final revision of the document.
- Carmen Camacho Leis: prehospital setting, final revision of the document.
- Salvador Espinosa Ramírez: prehospital setting, final revision of the document.
- José Manuel Fandiño Orgeira: critical reading, final revision of the document.
- Manuel José Vázquez Lima: critical reading, final revision of the document.
- Miguel Mayo-Yáñez: FONA contribution, final revision of the document.
- Pablo Parente-Arias: FONA contribution, final revision of the document.
- Jon Alexander Sistiaga-Suárez: critical reading, final revision of the document.
- Manuel Bernal-Sprekelsen: critical reading, final revision of the document.
- Pedro Charco-Mora: Coordination, Algorithm for unanticipated difficult tracheal intubation, draft ergonomic options, draft teaching and training, literature review, critical reading, final revision of the document

**Conflict of interests**

MAGR received lecture honoraria from Medtronic.

XOF received honoraria for lecture and practical workshop on neuromuscular block from Merck Sharp & Dohme.

RCF received honoraria for lectures from Fresenius Kabi.

AAG received honoraria for lectures from Merck Sharp & Dohme and 3 M Edwards.

## Delphi expert panel

Manuel Á. Gómez-Ríos, José Alfonso Sastre, Xavier Onrubia-Fuertes, Teresa López, Alfredo Abad-Gurumeta, José Carlos Garzón, Vicente Martínez-Pons, Marta Casalderrey-Rivas, Miguel Ángel Fernández-Vaquero, Eugenio Martínez-Hurtado, Ricardo Martín-Larrauri, Laura Reviriego-Agudo, Javier García-Fernández, Pedro Charco-Mora, Raquel García Álvarez, Alfredo Panadero Sánchez, Alejandra Prieto Gundín, María Luisa Santos Marqués, David Domínguez García, Irma, María Barrio, Uxío García-AlDAO, Aixa Espinosa, Carmen M. Holgado Pascual, Jesús Carazo Cordobés, Cristobal Añez Simón, Natividad Quesada Gimeno, Marina Gómez-Morán Quintana, Silvia Bermejo, Pilar Cabrerizo Torrente, Francisca Llobell, Roque J. Company Teuler, Teresa del Castillo Fdez de Betoño, Carlos González Perrino, and Paola Hurtado.

## External reviewers

Jaideep Pandit, Luis Gaitini, Tomasz Gaszyński, and Pavel Michalek

## Contributors

A list of contributors can be found in the supplementary material.

## Acknowledgements

We would like to thank Sandra Tejero Muñoz for illustrating figure 5 of this consensus document.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.redar.2023.08.001>.

## References

1. Artine CA, Roy S, Hagberg CA. The difficult airway. *Otolaryngol Clin North Am.* 2019;52(6):1115–25.
2. Collins SR, Blank RS. Fiberoptic intubation: an overview and update. *Respir Care.* 2014;59(6):865–78, discussion 878-880.
3. Aziz MF, Kristensen MS. From variance to guidance for awake tracheal intubation. *Anesthesia.* 2020;75(4):442–6.
4. Cabrera JL, Auerbach JS, Merelman AH, Levitan RM. The high-risk airway. *Emerg Med Clin North Am.* 2020;38(2):401–17.
5. Gómez-Ríos MA, Gaitini L, Matter I, Somri M. Guidelines and algorithms for managing the difficult airway. *Rev Esp Anestesiol Reanim.* 2018;65(1):41–8.
6. Gil K, Diemunsch P. Flexible scope intubation techniques. In: Hagberg CA, editor. *Benumof and Hagberg's Airway Management* (4th Edition). Philadelphia: W.B. Saunders; 2018. p. 428–570.
7. Wong P, Wong J, Mok MU. Anaesthetic management of acute airway obstruction. *Singapore Med J.* 2016;57(3):110–7.
8. Rosenblatt WH, Yanez ND. A decision tree approach to airway management pathways in the 2022 difficult airway algorithm of the American society of anesthesiologists. *Anesth Analg.* 2022;134(5):910–5.
9. Nekhendzy V, Biro P. Airway Management in Head and Neck Surgery. In: Hagberg CA, editor. *Benumof and Hagberg's Airway Management*. 4th edition Philadelphia: W.B. Saunders; 2018. p. 668–91, e5.
10. Hohn A, Kauliņš T, Hinkelbein J, Kauliņa K, Kopp A, Russo SG, et al. Awake tracheotomy in a patient with stridor and dyspnoea caused by a sizeable malignant thyroid tumor: a case report and short review of the literature. *Clin Case Rep.* 2017;5(11):1891–5.
11. Van Zundert AA, Endlich Y, Beckmann LA, Bradley WP, Chapman GA, Heard AM, et al. 2021 Update on airway management from the Anaesthesia Continuing Education Airway Management Special Interest Group. *Anaesth Intensive Care.* 2021;49(4):257–67.
12. Artine CA, Sanchez A. Preparation of the Patient for Awake Intubation. In: Hagberg CA, editor. *Benumof and Hagberg's Airway Management*. 4th edition Philadelphia: W.B. Saunders; 2018. p. 216–34, e4.
13. Vora J, Leslie D, Stacey M. Awake tracheal intubation. *BJA Educ.* 2022;22(8):298–305.
14. Law JA, Morris IR, Brousseau PA, de la Ronde S, Milne AD. The incidence, success rate, and complications of awake tracheal intubation in 1,554 patients over 12 years: an historical cohort study. *Can J Anaesth.* 2015;62(7):736–44.
15. Joseph TT, Gal JS, DeMaria S, Lin HM, Levine AI, Hyman JB. A retrospective study of success, failure, and time needed to perform awake intubation. *Anesthesiology.* 2016;125(1):105–14.
16. Cabrini L, Bairdo Redaelli M, Ball L, Filippini M, Fominskiy E, Pintaudi M, et al. Awake fiberoptic intubation protocols in the operating room for anticipated difficult airway: a systematic review and meta-analysis of randomized controlled trials. *Anesth Analg.* 2019;128(5):971–80.
17. Moore A, Schircker T. Awake videolaryngoscopy versus fiberoptic bronchoscopy. *Curr Opin Anaesthesiol.* 2019;32(6):764–8.
18. Bradley JA, Urman RD, Yao D. Challenging the traditional definition of a difficult intubation: what is difficult? *Anesth Analg.* 2019;128(3):584–6.
19. Leslie D, Stacey M. Awake intubation. *Continuing Education in Anaesthesia Critical Care & Pain.* 2014;15(2):64–7.
20. Badiger S, John M, Fearnley RA, Ahmad I. Optimizing oxygenation and intubation conditions during awake fibre-optic intubation using a high-flow nasal oxygen-delivery system. *Br J Anaesth.* 2015;115(4):629–32.
21. Ben-Menachem E, McKenzie J, O'Sullivan C, Havryk AP. High-flow nasal oxygen versus standard oxygen during flexible bronchoscopy in lung transplant patients: a randomized controlled trial. *J Bronchology Interv Pulmonol.* 2020;27(4):259–65.
22. Kim SH, Bang S, Lee KY, Park SW, Park JY, Lee HS, et al. Comparison of high flow nasal oxygen and conventional nasal cannula during gastrointestinal endoscopic sedation in the prone position: a randomized trial. *Can J Anaesth.* 2021;68(4):460–6.
23. Patel A, Nouraei SA. Transnasal Humidified Rapid-Insufflation Ventilatory Exchange (THRIVE): a physiological method of increasing apnoea time in patients with difficult airways. *Anaesthesia.* 2015;70(3):323–9.
24. Endlich Y, Hore PJ, Baker PA, Beckmann LA, Bradley WP, Chan KLE, et al. Updated guideline on equipment to manage difficult airways: Australian and New Zealand College of Anaesthetists. *Anaesth Intensive Care.* 2022;310057X221082664.
25. Zou T, Huang Z, Hu X, Cai G, He M, Wang S, et al. Clinical application of a novel endoscopic mask: a randomized controlled,

- multi-center trial in patients undergoing awake fiberoptic bronchoscopic intubation. *BMC Anesthesiol.* 2017;17(1):79.
26. Macfarlane AJR, Gitman M, Bornstein KJ, El-Boghdadly K, Weinberg G. Updates in our understanding of local anaesthetic systemic toxicity: a narrative review. *Anesthesia.* 2021;76 Suppl 1:27–39.
27. Dhooria S, Chaudhary S, Ram B, Sehgal IS, Muthu V, Prasad KT, et al. A randomized trial of nebulized lignocaine, lignocaine spray, or their combination for topical anesthesia during diagnostic flexible bronchoscopy. *Chest.* 2020;157(1):198–204.
28. McCambridge AJ, Boesch RP, Mullon JJ. Sedation in bronchoscopy: a review. *Clin Chest Med.* 2018;39(1):65–77.
29. Williams K, Barker G, Harwood R, Woodall N. Plasma lidocaine levels during local anaesthesia of the airway. *Anesthesia.* 2003;58(5):508–9.
30. Simmons ST, Schleich AR. Airway regional anesthesia for awake fiberoptic intubation. *Reg Anesth Pain Med.* 2002;27(2):180–92.
31. Li A, D'Costa J. Trans-cricoid thyroid injection of local anaesthesia: a serious complication. *BMJ Case Rep.* 2021;14(8).
32. Butler K, Winters M. The physiologically difficult intubation. *Emerg Med Clin North Am.* 2022;40(3):615–27.
33. Takita K, Morimoto Y, Kemmotsu O. Tracheal lidocaine attenuates the cardiovascular response to endotracheal intubation. *Can J Anaesth.* 2001;48(8):732–6.
34. Woodall NM, Harwood RJ, Barker GL. Complications of awake fibreoptic intubation without sedation in 200 healthy anaesthetists attending a training course. *Br J Anaesth.* 2008;100(6):850–5.
35. Johnston KD, Rai MR. Conscious sedation for awake fibreoptic intubation: a review of the literature. *Can J Anaesth.* 2013;60(6):584–99.
36. Murphy T, Howes B. Current practice for awake fibreoptic intubation - some unanswered questions. *Anesthesia.* 2017;72(6):678–81.
37. Wilson WM, Smith AF. The emerging role of awake videolaryngoscopy in airway management. *Anesthesia.* 2018;73(9):1058–61.
38. He XY, Cao JP, He Q, Shi XY. Dexmedetomidine for the management of awake fibreoptic intubation. *Cochrane Database Syst Rev.* 2014;(1):CD009798.
39. Zhou LJ, Fang XZ, Gao J, Zhangm Y, Tao LJ. Safety and efficacy of dexmedetomidine as a sedative agent for performing awake intubation: a meta-analysis. *Am J Ther.* 2016;23(6):e1788–800.
40. Tang ZH, Chen Q, Wang X, Su N, Xia Z, Wang Y, et al. A systematic review and meta-analysis of the safety and efficacy of remifentanil and dexmedetomidine for awake fiberoptic endoscope intubation. *Medicine (Baltimore).* 2021;100(14):e25324.
41. Vennila R, Hall A, Ali M, Bhuiyan N, Pirotta D, Raw DA. Remifentanil as single agent to facilitate awake fibreoptic intubation in the absence of premedication. *Anesthesia.* 2011;66(5):368–72.
42. Tsai CJ, Chu KS, Chen TI, Lu DV, Wang HM, Lu IC. A comparison of the effectiveness of dexmedetomidine versus propofol target-controlled infusion for sedation during fibreoptic nasotracheal intubation. *Anaesthesia.* 2010;65(3):254–9.
43. Jiang J, Ma DX, Li B, Wu AS, Xue FS. Videolaryngoscopy versus fiberoptic bronchoscope for awake intubation - a systematic review and meta-analysis of randomized controlled trials. *Ther Clin Risk Manag.* 2018;14:1955–63.
44. Benumof JL. Awake intubations are alive and well. *Can J Anaesth.* 2015;62(7):723–6.
45. El-Boghdadly K, Onwochei DN, Cuddihy J, Ahmad I. A prospective cohort study of awake fibreoptic intubation practice at a tertiary centre. *Anesthesia.* 2017;72(6):694–703.
46. Fitzgerald E, Hodzovic I, Smith AF. 'From darkness into light': time to make awake intubation with videolaryngoscopy the primary technique for an anticipated difficult airway? *Anesthesia.* 2015;70(4):387–92.
47. Alhomary M, Ramadan E, Curran E, Walsh SR. Videolaryngoscopy vs. fibreoptic bronchoscopy for awake tracheal intubation: a systematic review and meta-analysis. *Anesthesia.* 2018;73(9):1151–61.
48. Desai N, Ratnayake G, Onwochei DN, El-Boghdadly K, Ahmad I. Airway devices for awake tracheal intubation in adults: a systematic review and network meta-analysis. *Br J Anaesth.* 2021;127(4):636–47.
49. Ahmad I, Bailey CR. Time to abandon awake fibreoptic intubation? *Anesthesia.* 2016;71(1):12–6.
50. Johnson DM, From AM, Smith RB, From RP, Maktabi MA. Endoscopic study of mechanisms of failure of endotracheal tube advancement into the trachea during awake fiberoptic orotracheal intubation. *Anesthesiology.* 2005;102(5):910–4.
51. Asai T, Shingu K. Difficulty in advancing a tracheal tube over a fibreoptic bronchoscope: incidence, causes and solutions. *Br J Anaesth.* 2004;92(6):870–81.
52. Dutta K, Sriganesh K, Chakrabarti D, Pruthi N, Reddy M. Cervical spine movement during awake orotracheal intubation with fiberoptic scope and mcgrath videolaryngoscope in patients undergoing surgery for cervical spine instability: a randomized control trial. *J Neurosurg Anesthesiol.* 2020;32(3):249–55.
53. Cabrini L, Baiardo Redaelli M, Filippini M, Fominskij E, Pasin L, Pintaudi M, et al. Tracheal intubation in patients at risk for cervical spinal cord injury: a systematic review. *Acta Anaesthesiol Scand.* 2020;64(4):443–54.
54. Wiles MD. Airway management in patients with suspected or confirmed traumatic spinal cord injury: a narrative review of current evidence. *Anesthesia.* 2022;77(10):1120–8.
55. Chan JK, Ng I, Ang JP, Koh SM, Lee K, Mezzavia P, et al. Randomised controlled trial comparing the Ambu® aScope™2 with a conventional fibreoptic bronchoscope in orotracheal intubation of anaesthetised adult patients. *Anaesth Intensive Care.* 2015;43(4):479–84.
56. Krugel V, Bathory I, Frascarolo P, Schoettker P. Comparison of the single-use Ambu® aScope™ 2 vs the conventional fibrescope for tracheal intubation in patients with cervical spine immobilisation by a semirigid collar\*. *Anesthesia.* 2013;68(1):21–6.
57. Mouritsen JM, Ehlers L, Kovaleva J, Ahmad I, El-Boghdadly K. A systematic review and cost effectiveness analysis of reusable vs. single-use flexible bronchoscopes. *Anesthesia.* 2020;75(4):529–40.
58. Terjesen CL, Kovaleva J, Ehlers L. Early assessment of the likely cost effectiveness of single-use flexible video bronchoscopes. *Pharmacoecon Open.* 2017;1(2):133–41.
59. Gómez-Ríos MA, Nieto Serradilla L. Combined use of an Airtraq® optical laryngoscope, Airtraq video camera, Airtraq wireless monitor, and a fibreoptic bronchoscope after failed tracheal intubation. *Can J Anaesth.* 2011;58(4):411–2.
60. Mazzinari G, Rovira L, Henao L, Ortega J, Casasempere A, Fernandez Y, et al. Effect of dynamic versus stylet-guided intubation on first-attempt success in difficult airways undergoing glidescope laryngoscopy: a randomized controlled trial. *Anesth Analg.* 2019;128(6):1264–71.
61. Lenhardt R, Burkhardt MT, Brock GN, Kanchi-Kandadai S, Sharma R, Akça O. Is video laryngoscope-assisted flexible tracheoscope intubation feasible for patients with predicted difficult airway? A prospective, randomized clinical trial. *Anesth Analg.* 2014;118(6):1259–65.
62. Lim WY, Wong P. Awake supraglottic airway guided flexible bronchoscopic intubation in patients with anticipated difficult airways: a case series and narrative review. *Korean J Anesthesiol.* 2019;72(6):548–57.
63. Jadhav T, Sriganesh K, Reddy M, Chakrabarti D. Comparative study of fiberoptic guided versus intubating laryngeal

- mask airway assisted awake orotracheal intubation in patients with unstable cervical spine. *Minerva Anestesiol.* 2017;83(8):804–11.
64. Kristensen MS, McGuire B. Managing and securing the bleeding upper airway: a narrative review. *Can J Anaesth.* 2020;67(1):128–40.
65. Yan CL, Zhang YQ, Chen Y, Qv ZY, Zuo MZ. Comparison of SaCoVLM™ video laryngeal mask-guided intubation and i-gel combined with flexible bronchoscopy-guided intubation in airway management during general anesthesia: a non-inferiority study. *BMC Anesthesiol.* 2022;22(1):302.
66. Gómez-Ríos MA, López T, Sastre JA, Gaszyński T, Van Zundert AAJ. Video laryngeal masks in airway management. *Expert Rev Med Devices.* 2022;19(11):847–58.
67. Patel A, Pearce A. Progress in management of the obstructed airway. *Anaesthesia.* 2011;66 Suppl 2:93–100.
68. Rosenstock CV, Hodzovic I. Awake Tracheal Intubation. In: Kristensen MS, Cook T, editors. *Core Topics in Airway Management.* 3 ed. Cambridge: Cambridge University Press; 2020. p. 80–6.
69. Su K, Gao X, Xue FS, Ding GN, Zhang Y, Tian M. Difficult tracheal tube passage and subglottic airway injury during intubation with the GlideScope. *Anaesthesia.* 2017;72(4):504–11.
70. Barker KF, Bolton P, Cole S, Coe PA. Ease of laryngeal passage during fibreoptic intubation: a comparison of three endotracheal tubes. *Acta Anaesthesiol Scand.* 2001;45(5):624–6.
71. Hung KC, Chen JY, Feng IJ, Chiang MH, Wu SC, Chen IW, et al. Efficacy and airway complications of Parker Flex-Tip tubes and standard endotracheal tubes during airway manipulation: a meta-analysis and trial sequential analysis. *Eur J Anaesthesiol.* 2021;38(8):813–24.
72. Yamauchi H, Nakayama M, Yamamoto S, Sata M, Mato N, Bando M, et al. A comparative study of the Parker Flex-Tip tube versus standard portex tube for oral fiberoptic intubation in bronchoscopy performed by pulmonologists with limited experience. *Respir Investig.* 2021;59(2):223–7.
73. Sugiyama K, Manabe Y, Kohjitani A. The Parker Flex-Tip® tube prevents subglottic impingement on the tracheal wall during nasotracheal intubation. *Anesth Analg.* 2012;115(1):212–3.
74. Suzuki A, Ohmura T, Tampo A, Goto Y, Oikawa O, Kunisawa T, et al. Parker Flex-Tip Tube® provides higher intubation success with the Pentax-AWS Airwayscope® despite the AWS tip being inserted into the vallecula. *J Anesth.* 2012;26(4):614–6.
75. Greer JR, Smith SP, Strang T. A comparison of tracheal tube tip designs on the passage of an endotracheal tube during oral fiberoptic intubation. *Anesthesiology.* 2001;94(5):729–31.
76. Kristensen MS. The Parker Flex-Tip tube versus a standard tube for fiberoptic orotracheal intubation: a randomized double-blind study. *Anesthesiology.* 2003;98(2):354–8.
77. Radesic BP, Winkelman C, Einsporn R, Kless J. Ease of intubation with the Parker Flex-Tip or a standard Mallinckrodt endotracheal tube using a video laryngoscope (GlideScope). *AANA J.* 2012;80(5):363–72.
78. Jafari A, Gharaei B, Kamranmanesh MR, Aghamohammadi H, Nobahar MR, Poorzamany M, et al. Wire reinforced endotracheal tube compared with Parker Flex-Tip tube for oral fiberoptic intubation: a randomized clinical trial. *Minerva Anestesiol.* 2014;80(3):324–9.
79. Lomax SL, Johnston KD, Marfin AG, Yentis SM, Kathawaroo S, Popat MT. Nasotracheal fibreoptic intubation: a randomised controlled trial comparing the GlideRite® (Parker-Flex® Tip) nasal tracheal tube with a standard pre-rotated nasal RAE™ tracheal tube. *Anaesthesia.* 2011;66(3):180–4.
80. McNelis U, Syndercombe A, Harper I, Duggan J. The effect of cricoid pressure on intubation facilitated by the gum elastic bougie. *Anaesthesia.* 2007;62(5):456–9.
81. Koga K, Asai T, Latto IP, Vaughan RS. Effect of the size of a tracheal tube and the efficacy of the use of the laryngeal mask for fibrescope-aided tracheal intubation. *Anaesthesia.* 1997;52(2):131–5.
82. Greenland KB, Segal R, Acott C, Edwards MJ, Teoh WH, Bradley WP. Observations on the assessment and optimal use of video-laryngoscopes. *Anaesth Intensive Care.* 2012;40(4):622–30.
83. Karmali S, Rose P. Tracheal tube size in adults undergoing elective surgery - a narrative review. *Anaesthesia.* 2020;75(11):1529–39.
84. Farrow S, Farrow C, Soni N. Size matters: choosing the right tracheal tube. *Anaesthesia.* 2012;67(8):815–9.
85. Hillel AT, Karatayli-Ozgursoy S, Samad I, Best SR, Pandian V, Giraldez L, et al. Predictors of posterior glottic stenosis: a multi-institutional case-control study. *Ann Otol Rhinol Laryngol.* 2016;125(3):257–63.
86. Benjamin B. Prolonged intubation injuries of the larynx: endoscopic diagnosis, classification, and treatment. *Ann Otol Rhinol Laryngol.* 2018;127(8):492–507.
87. Sudhoff TH, Seidl RO, Estel B, Coordes A. Association of oversized tracheal tubes and cuff overinsufflation with postintubation tracheal ruptures. *Clin Exp Otorhinolaryngol.* 2015;8(4):409–15.
88. El-Boghdady K, Bailey CR, Wiles MD. Postoperative sore throat: a systematic review. *Anaesthesia.* 2016;71(6):706–17.
89. Hu B, Bao R, Wang X, Liu S, Tao T, Xie Q, et al. The size of endotracheal tube and sore throat after surgery: a systematic review and meta-analysis. *PLoS One.* 2013;8(10):e74467.
90. Esianor BI, Campbell BR, Casey JD, Du L, Wright A, Steitz B, et al. Endotracheal tube size in critically ill patients. *JAMA Otolaryngol Head Neck Surg.* 2022;148(9):849–53.
91. Cho HY, Yang SM, Jung CW, Cheun H, Lee HC, Park HP, et al. A randomised controlled trial of 7.5-mm and 7.0-mm tracheal tubes vs. 6.5-mm and 6.0-mm tracheal tubes for men and women during laparoscopic surgery. *Anaesthesia.* 2022;77(1):54–8.
92. Wirth S, Seywert L, Spaeth J, Schumann S. Compensating artificial airway resistance via active expiration assistance. *Respir Care.* 2016;61(12):1597–604.
93. Hwang JY, Park SH, Han SH, Park SJ, Park SK, Kim JH. The effect of tracheal tube size on air leak around the cuffs. *Korean J Anesthesiol.* 2011;61(1):24–9.
94. Brodsky MB, Akst LM, Jedlanek E, Pandian V, Blackford B, Price C, et al. Laryngeal injury and upper airway symptoms after endotracheal intubation during surgery: a systematic review and meta-analysis. *Anesth Analg.* 2021;132(4):1023–32.
95. Merelman AH, Perlmutter MC, Strayer RJ. Alternatives to rapid sequence intubation: contemporary airway management with ketamine. *West J Emerg Med.* 2019;20(3):466–71.
96. Morris IR. Preparation for Awake Intubation. In: Hung O, Murphy MF, editors. *Management of the Difficult and Failed Airway.* 3rd edition McGraw-Hill Education; 2017. p. 39–85.
97. Shukairy MK, Chadwick L, LaPorte CM, Pudwill J, Syslo JA, Fitzgerald J, et al. Implementing an interprofessional difficult airway response team to identify and manage high-risk airways. *Otolaryngol Head Neck Surg.* 2023.
98. Aziz S, Foster E, Lockey DJ, Christian MD. Emergency scalpel cricothyroidotomy use in a prehospital trauma service: a 20-year review. *Emerg Med J.* 2021;38(5):349–54.
99. Price TM, McCoy EP. Emergency front of neck access in airway management. *BJA Educ.* 2019;19(8):246–53.
100. Kwon YS, Lee CA, Park S, Ha SO, Sim YS, Baek MS. Incidence and outcomes of cricothyrotomy in the cannot intubate, cannot oxygenate situation. *Medicine (Baltimore).* 2019;98(42):e17713.
101. Bribriesco A, Patterson GA. Cricothyroid approach for emergency access to the airway. *Thorac Surg Clin.* 2018;28(3):435–40.

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

102. Kaufman MR, Alfonso KP, Burke K, Aouad RK. Awake vs sedated tracheostomies: a review and comparison at a single institution. *Otolaryngol Head Neck Surg.* 2018;159(5):830–4.
103. Ho AM, Chung DC, To EW, Karmakar MK. Total airway obstruction during local anesthesia in a non-sedated patient with a compromised airway. *Can J Anaesth.* 2004;51(8):838–41.
104. Pandian V, Ghazi TU, He MQ, Isak E, Saleem A, Semler LR, et al. Multidisciplinary difficult airway team characteristics, airway securement success, and clinical outcomes: a systematic review. *Ann Otol Rhinol Laryngol.* 2022;34894221123124.
105. Morris IR. Flexible Bronchoscopic Intubation. In: Hung O, Murphy MF, editors. *Management of the Difficult and Failed Airway.* 3rd edition McGraw-Hill Education; 2017. p. 172–97.
106. Ffrench-O'Carroll R, Fitzpatrick K, Jonker WR, Choo M, Tujjar O. Maintaining oxygenation with high-flow nasal cannula during emergent awake surgical tracheostomy. *Br J Anaesth.* 2017;118(6):954–5.
107. Adams TRP, Ricciardelli A. Airway fire during awake tracheostomy using high-flow nasal oxygen. *Anaesthetist Rep.* 2020;8(1):25–7.
108. O'Dell K. Predictors of difficult intubation and the otolaryngology perioperative consult. *Anesthesiol Clin.* 2015;33(2):279–90.
109. Sagiv D, Nachalon Y, Mansour J, Glikson E, Alon EE, Yakirevitch A, et al. Awake tracheostomy: indications, complications and outcome. *World J Surg.* 2018;42(9):2792–9.
110. Malpas G, Hung O, Gilchrist A, Wong C, Kent B, Hirsch GM, et al. The use of extracorporeal membrane oxygenation in the anticipated difficult airway: a case report and systematic review. *Can J Anaesth.* 2018;65(6):685–97.
111. Karim AS, Son AY, Suen R, Walter JM, Saine M, Kim SS, et al. Pre-intubation veno-venous extracorporeal membrane oxygenation in patients at risk for respiratory decompensation. *J Extra Corpor Technol.* 2020;52(1):52–7.
112. Hang D, Tawil JN, Fierro MA. Venovenous extracorporeal membrane oxygenation for rigid bronchoscopy and carinal tumor resection in decompensating patients. *Anesthesiology.* 2020;132(1):156.
113. Pu H, Huang X, Allingstrup MJ, Doig GS, Liang Z. Airway reconstruction supported by venovenous extracorporeal membrane oxygenation for patients with malignant critical central airway obstructions: a case series. *J Clin Anesth.* 2020;61:109690.
114. Gulkarov I, Khusid E, Worku B, Demissie S, Guerges M, Salemi A, et al. Meta-analysis of the effect of vascular complications on mortality in patients undergoing femoral venoarterial extracorporeal membrane oxygenation. *Ann Vasc Surg.* 2021;71:488–95.
115. Zangrillo A, Landoni G, Biondi-Zoccali G, Greco M, Greco T, Frati G, et al. A meta-analysis of complications and mortality of extracorporeal membrane oxygenation. *Crit Care Resusc.* 2013;15(3):172–8.
116. Yunoki K, Miyawaki I, Yamazaki K, Mima H. Extracorporeal membrane oxygenation-assisted airway management for difficult airways. *J Cardiothorac Vasc Anesth.* 2018; 32(6):2721–5.
117. Anton-Martin P, Bhattacharai P, Rycus P, Raman L, Potera R. The use of extracorporeal membrane oxygenation in life-threatening foreign body aspiration: case series, review of extracorporeal life support organization registry data, and systematic literature review. *J Emerg Med.* 2019;56(5):523–9.
118. Huitink JM, Cook T. The Epidemiology of Airway Management Complications. In: Kristensen MS, Cook T, editors. *Core Topics in Airway Management.* 3rd ed. Cambridge: Cambridge University Press; 2020. p. 22–37.
119. Chrimes N, Higgs A, Rehak A. Lost in transition: the challenges of getting airway clinicians to move from the upper airway to the neck during an airway crisis. *Br J Anaesth.* 2020;125(1):e38–46.
120. Myatra SN. Airway management in the critically ill. *Curr Opin Crit Care.* 2021;27(1):37–45.
121. Nørskov AK, Rosenstock CV, Wetterslev J, Astrup G, Afshari A, Lundstrøm LH. Diagnostic accuracy of anaesthesiologists' prediction of difficult airway management in daily clinical practice: a cohort study of 188 064 patients registered in the Danish Anaesthesia Database. *Anaesthesia.* 2015;70(3):272–81.
122. Natt BS, Malo J, Hypes CD, Sakles JC, Mosier JM. Strategies to improve first attempt success at intubation in critically ill patients. *Br J Anaesth.* 2016;117 Suppl 1:i60–8.
123. Hung O, Murphy MF. Context-Sensitive Airway Management. In: Hung O, Murphy MF, editors. *Management of the Difficult and Failed Airway.* 3rd edition McGraw-Hill Education; 2017. p. 136–42.
124. Crawley SM, McGuire B. New dimensions in airway management: risks for healthcare staff. *Anaesthesia.* 2020;75(11):1420–3.
125. Marshall SD, Pandit JJ. Radical evolution: the 2015 Difficult Airway Society guidelines for managing unanticipated difficult or failed tracheal intubation. *Anaesthesia.* 2016;71(2):131–7.
126. El-Boghdady K, Aziz MF. Face-mask ventilation: the neglected essentials? *Anaesthesia.* 2019;74(10):1227–30.
127. Chrimes N. The Vortex: a universal' high-acuity implementation tool' for emergency airway management. *Br J Anaesth.* 2016;117 Suppl 1:i20–7.
128. Liao CC, Liu FC, Li AH, Yu HP. Video laryngoscopy-assisted tracheal intubation in airway management. *Expert Rev Med Devices.* 2018;1–11.
129. Amalric M, Larcher R, Brunot V, Garnier F, De Jong A, Moulaire Rigollet V, et al. Impact of videolaryngoscopy expertise on first-attempt intubation success in critically ill patients. *Crit Care Med.* 2020;48(10):e889–96.
130. Bodily JB, Webb HR, Weiss SJ, Braude DA. Incidence and duration of continuously measured oxygen desaturation during emergency department intubation. *Ann Emerg Med.* 2016;67(3):389–95.
131. Kerslake D, Oglesby AJ, Di Rollo N, James E, McKeown DW, Ray DC, investigators E. Tracheal intubation in an urban emergency department in Scotland: a prospective, observational study of 3738 intubations. *Resuscitation.* 2015;89:20–4.
132. Goto T, Watase H, Morita H, Nagai H, Brown CA, Brown DF, et al., Investigators JEMN. Repeated attempts at tracheal intubation by a single intubator associated with decreased success rates in emergency departments: an analysis of a multicentre prospective observational study. *Emerg Med J.* 2015;32(10):781–6.
133. Kim J, Kim K, Kim T, Rhee JE, Jo YH, Lee JH, et al. The clinical significance of a failed initial intubation attempt during emergency department resuscitation of out-of-hospital cardiac arrest patients. *Resuscitation.* 2014;85(5):623–7.
134. Sakles JC, Chiu S, Mosier J, Walker C, Stoltz U. The importance of first pass success when performing orotracheal intubation in the emergency department. *Acad Emerg Med.* 2013;20(1):71–8.
135. Downey AW, Duggan LV, Adam Law J. A systematic review of meta-analyses comparing direct laryngoscopy with videolaryngoscopy. *Can J Anaesth.* 2021;68(5):706–14.
136. Hinkelbein J, Iovino I, De Robertis E, Kranke P. Outcomes in video laryngoscopy studies from 2007 to 2017: systematic review and analysis of primary and secondary endpoints for a core set of outcomes in video laryngoscopy research. *BMC Anesthesiol.* 2019;19(1):47.
137. Prekker ME, Driver BE, Trent SA, Resnick-Ault D, Seitz KP, Russell DW, et al. Video versus direct laryngoscopy for tracheal intubation of critically ill adults. *N Engl J Med.* 2023.
138. Russotto V, Lascarrou JB, Tassistro E, Parotto M, Antolini L, Bauer P, et al. Efficacy and adverse events profile of video-

- laryngoscopy in critically ill patients: subanalysis of the INTUBE study. *Br J Anaesth.* 2023.
139. Kim YS, Song J, Lim BG, Lee IO, Won YJ. Different classes of videoscopes and direct laryngoscopes for double-lumen tube intubation in thoracic surgery: a systematic review and network meta-analysis. *PLoS One.* 2020;15(8):e0238060.
140. Hoshijima H, Mihara T, Denawa Y, Ozaki M, Naya I, Shiga T, et al. Airtraq® is superior to the Macintosh laryngoscope for tracheal intubation: systematic review with trial sequential analysis. *Am J Emerg Med.* 2019;37(7):1367–8.
141. Jiang J, Ma DX, Li B, Wu AS, Xue FS. Videolaryngoscopy versus direct laryngoscopy for nasotracheal intubation: A systematic review and meta-analysis of randomised controlled trials. *J Clin Anesth.* 2019;52:6–16.
142. Arulkumaran N, Lowe J, Ions R, Mendoza M, Bennett V, Dunser MW. Videolaryngoscopy versus direct laryngoscopy for emergency orotracheal intubation outside the operating room: a systematic review and meta-analysis. *Br J Anaesth.* 2018;120(4):712–24.
143. Hoshijima H, Denawa Y, Tominaga A, Nakamura C, Shiga T, Nagasaka H. Videolaryngoscope versus Macintosh laryngoscope for tracheal intubation in adults with obesity: a systematic review and meta-analysis. *J Clin Anesth.* 2018;44:69–75.
144. Liu TT, Li L, Wan L, Zhang CH, Yao WL. Videolaryngoscopy vs. Macintosh laryngoscopy for double-lumen tube intubation in thoracic surgery: a systematic review and meta-analysis. *Anesthesia.* 2018;73(8):997–1007.
145. Jiang J, Ma D, Li B, Yue Y, Xue F. Video laryngoscopy does not improve the intubation outcomes in emergency and critical patients - a systematic review and meta-analysis of randomized controlled trials. *Crit Care.* 2017;21(1):288.
146. Pieters BMA, Maas EHA, Knape JTA, van Zundert AAJ. Videolaryngoscopy vs. direct laryngoscopy use by experienced anaesthetists in patients with known difficult airways: a systematic review and meta-analysis. *Anesthesia.* 2017;72(12):1532–41.
147. De Jong A, Molinari N, Conseil M, Coisel Y, Pouzeratte Y, Belafia F, et al. Video laryngoscopy versus direct laryngoscopy for orotracheal intubation in the intensive care unit: a systematic review and meta-analysis. *Intensive Care Med.* 2014;40(5):629–39.
148. Griesdale DE, Liu D, McKinney J, Choi PT. Glidescope® videolaryngoscopy versus direct laryngoscopy for endotracheal intubation: a systematic review and meta-analysis. *Can J Anaesth.* 2012;59(1):41–52.
149. Lu Y, Jiang H, Zhu YS. Airtraq laryngoscope versus conventional Macintosh laryngoscope: a systematic review and meta-analysis. *Anesthesia.* 2011;66(12):1160–7.
150. Hansel J, Rogers AM, Lewis SR, Cook TM, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adults undergoing tracheal intubation. *Cochrane Database Syst Rev.* 2022;4:CD011136.
151. Howle R, Onwochei D, Harrison SL, Desai N. Comparison of videolaryngoscopy and direct laryngoscopy for tracheal intubation in obstetrics: a mixed-methods systematic review and meta-analysis. *Can J Anaesth.* 2021;68(4):546–65.
152. Hoshijima H, Mihara T, Maruyama K, Denawa Y, Mizuta K, Shiga T, et al. C-MAC videolaryngoscope versus Macintosh laryngoscope for tracheal intubation: A systematic review and meta-analysis with trial sequential analysis. *J Clin Anesth.* 2018;49:53–62.
153. Hoshijima H, Mihara T, Maruyama K, Denawa Y, Takahashi M, Shiga T, et al. McGrath videolaryngoscope versus Macintosh laryngoscope for tracheal intubation: A systematic review and meta-analysis with trial sequential analysis. *J Clin Anesth.* 2018;46:25–32.
154. Huang HB, Peng JM, Xu B, Liu GY, Du B. Video laryngoscopy for endotracheal intubation of critically ill adults: a systemic review and meta-analysis. *Chest.* 2017;152(3):510–7.
155. Zhao BC, Huang TY, Liu KX. Video laryngoscopy for ICU intubation: a meta-analysis of randomised trials. *Intensive Care Med.* 2017;43(6):947–8.
156. Lewis SR, Butler AR, Parker J, Cook TM, Schofield-Robinson OJ, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation: a cochrane systematic review. *Br J Anaesth.* 2017;119(3):369–83.
157. Lewis SR, Butler AR, Parker J, Cook TM, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation. *Cochrane Database Syst Rev.* 2016;11:CD011136.
158. Hoshijima H, Kuratani N, Hirabayashi Y, Takeuchi R, Shiga T, Masaki E. Pentax Airway Scope® vs Macintosh laryngoscope for tracheal intubation in adult patients: a systematic review and meta-analysis. *Anesthesia.* 2014;69(8):911–8.
159. Bhattacharjee S, Maitra S, Baidya DK. A comparison between video laryngoscopy and direct laryngoscopy for endotracheal intubation in the emergency department: A meta-analysis of randomized controlled trials. *J Clin Anesth.* 2018;47:21–6.
160. Hoshijima H, Maruyama K, Mihara T, Mieda T, Shiga T, Nagasaka H. Airtraq® reduces the hemodynamic response to tracheal intubation using single-lumen tubes in adults compared with the Macintosh laryngoscope: A systematic review and meta-analysis of randomized control trials. *J Clin Anesth.* 2018;47:86–94.
161. Rombey T, Schieren M, Pieper D. Video versus direct laryngoscopy for inpatient emergency intubation in adults. *Dtsch Arztbl Int.* 2018;115(26):437–44.
162. Jiang J, Kang N, Li B, Wu AS, Xue FS. Comparison of adverse events between video and direct laryngoscopes for tracheal intubations in emergency department and ICU patients-a systematic review and meta-analysis. *Scand J Trauma Resusc Emerg Med.* 2020;28(1):10.
163. Wong P, Lim WY. Aligning difficult airway guidelines with the anesthetic COVID-19 guidelines to develop a COVID-19 difficult airway strategy: a narrative review. *J Anesth.* 2020;34(6):924–43.
164. De Jong A, Pardo E, Rolle A, Bodin-Lario S, Pouzeratte Y, Jaber S. Airway management for COVID-19: a move towards universal videolaryngoscope? *Lancet Respir Med.* 2020;8(6):555.
165. Gómez-Ríos M, Casans-Francés R, Abad-Gurumeta A, Esquinias A. The role of videolaryngoscopy in airway management of COVID-19 patients. *Anaesthesiol Intensive Ther.* 2020;52(4):344–5.
166. Hemmerling TM, Zaouter C. Videolaryngoscopy: is there a path to becoming a standard of care for intubation? *Anesth Analg.* 2020;131(4):1313–6.
167. De Jong A, Myatra SN, Roca O, Jaber S. How to improve intubation in the intensive care unit. Update on knowledge and devices. *Intensive Care Med.* 2022;48(10):1287–98.
168. Zhang J, Jiang W, Urdaneta F. Economic analysis of the use of video laryngoscopy versus direct laryngoscopy in the surgical setting. *J Comp Eff Res.* 2021;10(10):831–44.
169. Alsumali A, Noppens R. Cost effectiveness of video laryngoscopy for routine use in the operating room. *Trends in Anaesthesia and Critical Care.* 2018;23:10.
170. Samuels JD, Tangel VE, Lui B, Turnbull ZA, Pryor KO, White RS, et al. Adoption of video laryngoscopy by a major academic anesthesia department. *J Comp Eff Res.* 2021;10(2):101–8.
171. Theiler L, Cook T, Aziz M. Videolaryngoscopy. In: Kristensen MS, Cook T, editors. *Core Topics in Airway Management.* 3 ed. Cambridge: Cambridge University Press; 2020. p. 153–60.
172. McNarry AF, Patel A. The evolution of airway management - new concepts and conflicts with traditional practice. *Br J Anaesth.* 2017;119 suppl.1:i154–66.

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

173. Jaber S, De Jong A, Pelosi P, Cabrini L, Reignier J, Lascarrou JB. Videolaryngoscopy in critically ill patients. *Crit Care*. 2019;23(1):221.
174. Gomez-Rios MA, Sastre-Rincon JA, Mariscal-Flores M. Is direct laryngoscopy dead? Long live the video laryngoscopy. *Rev Esp Anestesiol Reanim*. 2019;66(4):177–80.
175. Natt B, Mosier J. Airway management in the critically ill patient. *Curr Anesthesiol Rep*. 2021;1:1–12.
176. De Jong A, Sfara T, Pouzeratte Y, Pensier J, Rolle A, Chanques G, et al. Videolaryngoscopy as a first-intention technique for tracheal intubation in unselected surgical patients: a before and after observational study. *Br J Anaesth*. 2022;129(4):624–34.
177. Dey S, Pradhan D, Saikia P, Bhattacharyya P, Khandelwal H, Adarsha KN. Intubation in the intensive care unit: C-MAC video laryngoscope versus Macintosh laryngoscope. *Med Intensiva*. 2020;44(3):135–41.
178. Schroeder RA, Pollard R, Dhakal I, Cooter M, Aronson S, Grichnik K, et al. Temporal trends in difficult and failed tracheal intubation in a regional community anesthetic practice. *Anesthesiology*. 2018;128(3):502–10.
179. Aziz MF, Brambrink AM, Healy DW, Willett AW, Shanks A, Tremper T, et al. Success of intubation rescue techniques after failed direct laryngoscopy in adults: a retrospective comparative analysis from the multicenter perioperative outcomes group. *Anesthesiology*. 2016;125(4):656–66.
180. Jayaraj AK, Siddiqui N, Abdelghany SMO, Balki M. Management of difficult and failed intubation in the general surgical population: a historical cohort study in a tertiary care centre. *Can J Anaesth*. 2022;69(4):427–37.
181. Berkow LC, Morey TE, Urdaneta F. The Technology of Video Laryngoscopy. *Anesth Analg*. 2018;126(5):1527–34.
182. Niforopoulou P, Pantazopoulos I, Demesticha T, Koudouna E, Xanthos T. Video-laryngoscopes in the adult airway management: a topical review of the literature. *Acta Anaesthesiol Scand*. 2010;54(9):1050–61.
183. Xue FS, He N, Liu JH, Liao X, Xu XZ, Zhang YM. More maneuvers to facilitate endotracheal intubation using the Airtraq laryngoscope in children with difficult airways. *Paediatr Anaesth*. 2009;19(9):916–8.
184. Ho AM, Ho AK, Mizubuti GB. Tracheal Intubation: The Proof is in the Bevel. *J Emerg Med*. 2018;55(6):821–6.
185. Grape S, Schoettker P. The role of tracheal tube introducers and stylets in current airway management. *J Clin Monit Comput*. 2017;31(3):531–7.
186. Shah A, Durnford K, Knecht L, Jacobson C, Runnels ST. A Consecutive Case Series of Rescue Intubations With the Articulating Total Control Introducer for Precision Tracheal Access. *A A Pract*. 2021;15(3):e01418.
187. Latimer AJ, Harrington B, Counts CR, Ruark K, Maynard C, Watase T, et al. Routine use of a bougie improves first-attempt intubation success in the out-of-hospital setting. *Ann Emerg Med*. 2021;77(3):296–304.
188. Driver BE, Prekker ME, Klein LR, Reardon RF, Miner JR, Fagerstrom ET, et al. Effect of use of a Bougie vs Endotracheal tube and stylet on first-attempt intubation success among patients with difficult airways undergoing emergency intubation: a randomized clinical trial. *JAMA*. 2018;319(21):2179–89.
189. Driver B, Dodd K, Klein LR, Buckley R, Robinson A, McGill JW, et al. The Bougie and first-pass success in the emergency department. *Ann Emerg Med*. 2017;70(4):473–8, e1.
190. Martin LD, Mhyre JM, Shanks AM, Tremper KK, Khetpal S. 3,423 emergency tracheal intubations at a university hospital: airway outcomes and complications. *Anesthesiology*. 2011;114(1):42–8.
191. Laurin EG. Endotracheal tube introducers (gum elastic bougie) for emergency intubation. *UpToDate*. Retrieved November 2021.
192. Russotto V, Myatra SN, Laffey JG. What's new in airway management of the critically ill. *Intensive Care Med*. 2019;45(11):1615–8.
193. Oxenham O, Pairaudeau C, Moody T, Mendonca C. Standard and flexible tip bougie for tracheal intubation using a non-channelled hyperangulated videolaryngoscope: a randomised comparison. *Anaesthesia*. 2022;77(12):1368–75.
194. Ruetzler K, Smereka J, Abelairas-Gomez C, Frass M, Dabrowski M, Bialka S, et al. Comparison of the new flexible tip bougie catheter and standard bougie stylet for tracheal intubation by anesthesiologists in different difficult airway scenarios: a randomized crossover trial. *BMC Anesthesiol*. 2020;20(1):90.
195. Heidegger T, Gerig HJ, Ulrich B, Schnider TW. Structure and process quality illustrated by fibreoptic intubation: analysis of 1612 cases. *Anaesthesia*. 2003;58(8):734–9.
196. Pandit JJ, Dravid RM, Iyer R, Popat MT. Orotracheal fibreoptic intubation for rapid sequence induction of anaesthesia. *Anaesthesia*. 2002;57(2):123–7.
197. Ching YH, Karlinoski RA, Chen H, Camporesi EM, Shah VV, Padhya TA, et al. Lingual traction to facilitate fiber-optic intubation of difficult airways: a single-anesthesiologist randomized trial. *J Anesth*. 2015;29(2):263–8.
198. Han SH, Oh AY, Jung CW, Park SJ, Kim JH, Nahm FS. The effect of the jaw-thrust manoeuvre on the ability to advance a tracheal tube over a bronchoscope during oral fibreoptic intubation. *Anaesthesia*. 2013;68(5):472–7.
199. Durga VK, Millns JP, Smith JE. Manoeuvres used to clear the airway during fibreoptic intubation. *Br J Anaesth*. 2001;87(2):207–11.
200. Adachi YU, Satomoto M, Higuchi H. Fiberoptic orotracheal intubation in the left semilateral position. *Anesth Analg*. 2002;94(2):477–8.
201. Wong DT, Mehta A, Tam AD, Yau B, Wong J. A survey of Canadian anesthesiologists' preferences in difficult intubation and cannot intubate, cannot ventilate situations. *Can J Anaesth*. 2014;61(8):717–26.
202. Orebaugh S, Snyder JV. Direct laryngoscopy and endotracheal intubation in adults. *UpToDate*. Retrieved November 2021.
203. Whitaker DK, Benson JP. Capnography standards for outside the operating room. *Curr Opin Anaesthesiol*. 2016;29(4): 485–92.
204. Russotto V, Myatra SN, Laffey JG, Tassistro E, Antolini L, Bauer P, et al. Intubation practices and adverse peri-intubation events in critically ill patients from 29 countries. *JAMA*. 2021;325(12):1164–72.
205. Straker TU, Felipe. Confirmation of Tracheal Intubation. In: Hagberg CA, editor. *Benumof and Hagberg's Airway Management*. 4th edition Philadelphia: W.B. Saunders; 2018. p. 540–50, e3.
206. Li J. Capnography alone is imperfect for endotracheal tube placement confirmation during emergency intubation. *J Emerg Med*. 2001;20(3):223–9.
207. Perkins GD, Graesner JT, Semeraro F, Olasveengen T, Soar J, Lott C, et al. European resuscitation council guidelines 2021: executive summary. *Resuscitation*. 2021;161:1–60.
208. Panchal AR, Bartos JA, Cabañas JG, Donnino MW, Drennan IR, Hirsch KG, et al. Part 3: adult basic and advanced life support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142 16\_suppl\_2:S366–468.
209. Ahmed A, Azim A. Difficult tracheal intubation in critically ill. *J Intensive Care*. 2018;6:49.
210. Jaber S, Jung B, Corne P, Sebbane M, Muller L, Chanques G, et al. An intervention to decrease complications related to endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Intensive Care Med*. 2010;36(2):248–55.
211. Russotto V, Cook TM. Capnography use in the critical care setting: why do clinicians fail to implement

- this safety measure? *Br J Anaesth.* 2021;127(5):661–4.
212. Sandroni C, De Santis P, D'Arrigo S. Capnography during cardiac arrest. *Resuscitation.* 2018;132:73–7.
213. Klein AA, Meek T, Allcock E, Cook TM, Mincher N, Morris C, et al. Recommendations for standards of monitoring during anaesthesia and recovery 2021: guideline from the Association of Anaesthetists. *Anaesthesia.* 2021;76(9):1212–23.
214. Kodali BS. Capnography outside the operating rooms. *Anesthesiology.* 2013;118(1):192–201.
215. Aminiahidashti H, Shafee S, Zamani Kiasari A, Sazgar M. Applications of end-tidal carbon Dioxide (ETCO<sub>2</sub>) monitoring in emergency department; a narrative review. *Emerg (Tehran).* 2018;6(1):e5.
216. Chrimes N, Higgs A, Hagberg CA, Baker PA, Cooper RM, Greif R, et al. Preventing unrecognised oesophageal intubation: a consensus guideline from the Project for Universal Management of Airways and international airway societies. *Anaesthesia.* 2022;77(12):1395–415.
217. Cook TM, Harrop-Griffiths W. Capnography prevents avoidable deaths. *BMJ.* 2019;364:l439.
218. Kerslake I, Kelly F. Uses of capnography in the critical care unit. *BJA Education.* 2017;17(5):178–83.
219. Pandit JJ, Young P, Davies M. Why does oesophageal intubation still go unrecognised? Lessons for prevention from the coroner's court. *Anaesthesia.* 2022;77(2):123–8.
220. Sahu AK, Bhoi S, Aggarwal P, Mathew R, Nayer J, et al. Endotracheal Tube Placement Confirmation by Ultrasonography: A Systematic Review and Meta-Analysis of more than 2500 Patients. *J Emerg Med.* 2020;59(2):254–64.
221. Gottlieb M, Holladay D, Peksa GD. Ultrasonography for the confirmation of endotracheal tube intubation: a systematic review and meta-analysis. *Ann Emerg Med.* 2018;72(6):627–36.
222. You-Ten KE, Siddiqui N, Teoh WH, Kristensen MS. Point-of-care ultrasound (POCUS) of the upper airway. *Can J Anaesth.* 2018;65(4):473–84.
223. Hansel J, Law JA, Chrimes N, Higgs A, Cook TM. Clinical tests for confirming tracheal intubation or excluding oesophageal intubation: a diagnostic test accuracy systematic review and meta-analysis. *Anaesthesia.* 2023;78(8):1020–30.
224. Salem MR. Verification of endotracheal tube position. *Anesthesiol Clin North Am.* 2001;19(4):813–39.
225. Jafferji D, Morris R, Levy N. Reducing the risk of confirmation bias in unrecognised oesophageal intubation. *Br J Anaesth.* 2019;122(4):e66–8.
226. Whitaker DK. Time for capnography - everywhere. *Anesthesia.* 2011;66(7):544–9.
227. Nassar BS, Schmidt GA. Capnography during critical illness. *Chest.* 2016;149(2):576–85.
228. Branson RD, Rodriguez D. Monitoring during transport. *Respir Care.* 2020;65(6):882–93.
229. Kreit JW. Volume capnography in the intensive care unit: potential clinical applications. *Ann Am Thorac Soc.* 2019;16(4):409–20.
230. Lam T, Nagappa M, Wong J, Singh M, Wong D, Chung F. Continuous pulse oximetry and capnography monitoring for postoperative respiratory depression and adverse events: a systematic review and meta-analysis. *Anesth Analg.* 2017;125(6):2019–29.
231. Kremeyer P, Böhm SH, Tusman G. Clinical use of volumetric capnography in mechanically ventilated patients. *J Clin Monit Comput.* 2020;34(1):7–16.
232. Schmidt GA. Monitoring gas exchange. *Respir Care.* 2020;65(6):729–38.
233. Budde AM, Kadar RB, Jabaley CS. Airway misadventures in adult critical care: a concise narrative review of managing lost or compromised artificial airways. *Curr Opin Anesthesiol.* 2022;35(2):130–6.
234. Cook TM, Woodall N, Harper J, Benger J. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 2: intensive care and emergency departments. *Br J Anaesth.* 2011;106(5):632–42.
235. Wang CH, Lee AF, Chang WT, Huang CH, Tsai MS, Chou E, et al. Comparing Effectiveness of Initial Airway Interventions for Out-of-Hospital Cardiac Arrest: A Systematic Review and Network Meta-analysis of Clinical Controlled Trials. *Ann Emerg Med.* 2020;75(5):627–36.
236. Wang H.E., Schmicker R.H., Daya M.R., Stephens S.W., Idris A.H., Carlson J.N., Coletta M.R., Herren H., Hansen M., Richmond N.J. et al. Effect of a Strategy of Initial Laryngeal Tube Insertion vs Endotracheal Intubation on 72-Hour Survival in Adults With Out-of-Hospital Cardiac Arrest: A Randomized Clinical Trial. *JAMA.* Volume 3202018. 769–778.
237. Benger JR, Kirby K, Black S, Brett SJ, Clout M, Lazaroo MJ, et al. Effect of a Strategy of a Supraglottic Airway Device vs Tracheal Intubation During Out-of-Hospital Cardiac Arrest on Functional Outcome: The AIRWAYS-2 Randomized Clinical Trial. *JAMA.* 2018;320(8):779–91.
238. Laurin EG. Extraluminal devices for emergency airway management in adults. *UpToDate.* Retrieved November 2021.
239. Timmermann A. Supraglottic airways in difficult airway management: successes, failures, use and misuse. *Anaesthesia.* 2011;66 Suppl 2:45–56.
240. Gordon J, Cooper RM, Parotto M. Supraglottic airway devices: indications, contraindications and management. *Minerva Anestesiol.* 2018;84(3):389–97.
241. Ramachandran SK, Kumar AM. Supraglottic airway devices. *Respir Care.* 2014;59(6):920–31.
242. Driver BE, Martel M, Lai T, Marko TA, Reardon RF. Use of the intubating laryngeal mask airway in the emergency department: A ten-year retrospective review. *Am J Emerg Med.* 2020;38(7):1367–72.
243. Lee DH, Stang J, Reardon RF, Martel ML, Driver BE, Braude DA. Rapid Sequence Airway with the Intubating Laryngeal Mask in the Emergency Department. *J Emerg Med.* 2021;61(5):550–7.
244. Lai CJ, Yeh YC, Tu YK, Cheng YJ, Liu CM, Fan SZ. Comparison of the efficacy of supraglottic airway devices in low-risk adult patients: a network meta-analysis and systematic review. *Sci Rep.* 2021;11(1):15074.
245. Cook TM, Kelly FE. Time to abandon the 'vintage' laryngeal mask airway and adopt second-generation supraglottic airway devices as first choice. *Br J Anaesth.* 2015;115(4):497–9.
246. Van Zundert AA, Kumar CM, Van Zundert TC. Malpositioning of supraglottic airway devices: preventive and corrective strategies. *Br J Anaesth.* 2016;116:579–82.
247. Hamaekers AE, Henderson JJ. Equipment and strategies for emergency tracheal access in the adult patient. *Anaesthesia.* 2011;66 Suppl 2:65–80.
248. Lønvik MP, Elden OE, Lunde MJ, Nordseth T, Bakkelund KE, Uleberg O. A prospective observational study comparing two supraglottic airway devices in out-of-hospital cardiac arrest. *BMC Emerg Med.* 2021;21(1):51.
249. Goldmann K, Hechtfischer C, Malik A, Kussin A, Freisburger C. Use of ProSeal laryngeal mask airway in 2114 adult patients: a prospective study. *Anesth Analg.* 2008;107(6):1856–61.
250. Koo CH, Oh AY, Jeon YT, Hwang JW, Ryu JH. Standard digit-based versus 90° rotation technique for supraglottic airway device insertion: a meta-analysis of randomized controlled trials. *Korean J Anesthesiol.* 2022;75(3):266–75.
251. Michalek P, Donaldson W, Vobrubova E, Hakl M. Complications associated with the use of supraglottic airway devices in perioperative medicine. *Biomed Res Int.* 2015;2015:746560.
252. Park JH, Lee JS, Nam SB, Ju JW, Kim MS. Standard versus rotation technique for insertion of supraglottic airway

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

- devices: systematic review and meta-analysis. *Yonsei Med J.* 2016;57(4):987–97.
253. Huh H, Cho JE, Lee SW, Kim HC. The effects of two-handed jaw thrust on i-gel™ placement in anesthetized non-paralyzed patients. *Minerva Anestesiol.* 2021;87(10):1109–16.
254. Baran Akkuş İ, Kavak Akelma F, Emlek M, Özkan D, Ergil J, Polat R. Comparison of the standard and triple airway maneuvering techniques for i-gel™ placement in patients undergoing elective surgery: a randomized controlled study. *J Anesth.* 2020;34(4):512–8.
255. Van Zundert AAJ, Gatt SP, Kumar CM, Van Zundert T, Pandit JJ. 'Failed supraglottic airway': an algorithm for suboptimally placed supraglottic airway devices based on videolaryngoscopy. *Br J Anaesth.* 2017;118(5):645–9.
256. Van Zundert AAJ, Gatt SP, Kumar CM, Van Zundert TCRV. Vision-guided placement of supraglottic airway device prevents airway obstruction: a prospective audit. *Br J Anaesth.* 2017;118(3):462–3.
257. van Zundert AAJ, Wyssusek KH, Pelecanos A, Roets M, Kumar CM. A prospective randomized comparison of airway seal using the novel vision-guided insertion of LMA-Supreme® and LMA-Protector®. *J Clin Monit Comput.* 2020;34(2):285–94.
258. Zhao L, Zhang J, Zhou Q, Jiang W. Comparison of a new visual stylet (Discopo)-guided laryngeal mask airway placement vs conventional blind technique: a prospective randomized study. *J Clin Anesth.* 2016;35:85–9.
259. Van Zundert AAJ, Kumar CM, Van Zundert TCRV, Gatt SP, Pandit JJ. The case for a 3rd generation supraglottic airway device facilitating direct vision placement. *J Clin Monit Comput.* 2021;35(2):217–24.
260. Gómez-Ríos MA, Freire-Vila E, Casans-Francés R, Pita-Fernández S. The Totaltrack TM video laryngeal mask: an evaluation in 300 patients. *Anesthesia.* 2019;74(6):751–7.
261. Gomez-Rios MA, Casans-Francés R, Freire-Vila E, Sastre JA, Lopez T, Garzon JC. A prospective evaluation of the Totaltrack video laryngeal mask in paralyzed, anesthetized obese patients. *J Clin Anesth.* 2020;61:109688.
262. Gómez-Ríos M, Freire-Vila E, Vizcaíno-Martínez L, Estévez-González E. The Totaltrack™: an initial evaluation. *Br J Anaesth.* 2015;115(5):798–9.
263. Gomez-Rios MA, Freire-Vila E, Calvo-Vecino JM. Use of the Totaltrack VLM as a rescue device following failed tracheal intubation. *Eur J Anaesthesiol.* 2019;36(3):237–9.
264. Gomez-Rios MA, Freire-Vila E, Abad-Gurumeta A, Barreto-Calvo P, Calvo-Vecino JM. Use of Totaltrack VLM as a rescue device after failed ventilation and tracheal intubation with LMA Fastrach in emergent difficult airways. *J Clin Anesth.* 2019;52:29–30.
265. Yoo S, Park SK, Kim WH, Hur M, Bahk JH, Lim YJ, et al. The effect of neck extension on success rate of blind intubation through Ambu® AuraGain™ laryngeal mask: a randomized clinical trial. *Can J Anaesth.* 2019;66(6):639–47.
266. Gomez-Rios MA, Bonome C. The totaltrack VLM: a novel video-assisted intubating laryngeal mask. *Minerva Anestesiol.* 2018;84(1):126–7.
267. Ahn E, Choi G, Kang H, Baek C, Jung Y, Woo Y, et al. Supraglottic airway devices as a strategy for unassisted tracheal intubation: A network meta-analysis. *PLoS One.* 2018;13(11):e0206804.
268. Liu EH, Goy RW, Chen FG. An evaluation of poor LMA CTrach views with a fiberoptic laryngoscope and the effectiveness of corrective measures. *Br J Anaesth.* 2006;97(6):878–82.
269. Nagata T, Kishi Y, Tanigami H, Hiuge Y, Sonoda S, Ohashi Y, et al. Oral gastric tube-guided insertion of the ProSeal™ laryngeal mask is an easy and noninvasive method for less experienced users. *J Anesth.* 2012;26(4):531–5.
270. van Zundert AAJ, Wyssusek KH. Epiglottis folding double with supraglottic airway devices. *Br J Anaesth.* 2018;120(4):884–5.
271. Fei M, Blair JL, Rice MJ, Edwards DA, Liang Y, Pilla MA, et al. Comparison of effectiveness of two commonly used two-handed mask ventilation techniques on unconscious apnoeic obese adults. *Br J Anaesth.* 2017;118(4):618–24.
272. Sato Y, Ikeda A, Ishikawa T, Isono S. How can we improve mask ventilation in patients with obstructive sleep apnea during anesthesia induction? *J Anesth.* 2013;27(1):152–6.
273. Joffe AM, Hetzel S, Liew EC. A two-handed jaw-thrust technique is superior to the one-handed ÈC-clamptechnique for mask ventilation in the apneic unconscious person. *Anesthesiology.* 2010;113(4):873–9.
274. Isono S. One hand, two hands, or no hands for maximizing airway maneuvers? *Anesthesiology.* 2008;109(4):576–7.
275. Matico AA. An anesthesiologist's perspective on the history of basic airway management: the modernera, 1960 to present. *Anesthesiology.* 2019;130(5):686–711.
276. Soltész S, Alm P, Mathes A, Hellmich M, Hinkelbein J. The effect of neuromuscular blockade on the efficiency of face-mask ventilation in patients difficult to facemask ventilate: a prospective trial. *Anesthesia.* 2017;72(12):1484–90.
277. Joffe AM, Ramaiah R, Donahue E, Galgon RE, Thilen SR, Spiekerman CF, et al. Ventilation by mask before and after the administration of neuromuscular blockade: a pragmatic non-inferiority trial. *BMC Anesthesiol.* 2015;15:134.
278. Sachdeva R, Kannan TR, Mendonca C, Patteril M. Evaluation of changes in tidal volume during mask ventilation following administration of neuromuscular blocking drugs. *Anesthesia.* 2014;69(8):826–31.
279. Patel A. Facemask ventilation before or after neuromuscular blocking drugs: where are we now? *Anesthesia.* 2014;69(8):811–5.
280. Cooper RM. Strengths and limitations of airway techniques. *Anesthesiol Clin.* 2015;33(2):241–55.
281. Mosier JM, Sakles JC, Law JA, Brown CA, Brindley PG. Tracheal Intubation in the Critically Ill. Where We Came from and Where We Should Go. *Am J Respir Crit Care Med.* 2020;201(7):775–88.
282. Wenzel V, Idris AH, Dörge V, Nolan JP, Parr MJ, Gabrielli A, et al. The respiratory system during resuscitation: a review of the history, risk of infection during assisted ventilation, respiratory mechanics, and ventilation strategies for patients with an unprotected airway. *Resuscitation.* 2001;49(2):123–34.
283. Bouvet L, Albert ML, Augris C, Boselli E, Ecochard R, Rabilloud M, et al. Real-time detection of gastric insufflation related to facemask pressure-controlled ventilation using ultrasonography of the antrum and epigastric auscultation in nonparalyzed patients: a prospective, randomized, double-blind study. *Anesthesiology.* 2014;120(2):326–34.
284. Bell D. Avoiding adverse outcomes when faced with 'difficulty with ventilation'. *Anesthesia.* 2003;58(10):945–8.
285. DuCanto J, Matico A. Noninvasive management of the airway. In: Hagberg C, Artine C, Aziz M, editors. *Hagberg and Benumof's Airway management.* 4th edition Philadelphia: Elsevier; 2018. p. 309–27.
286. Chrimes N, Cook TM. Critical airways, critical language. *Br J Anaesth.* 2017;118(5):649–54.
287. Cook TM, Woodall N, Frerk C. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 1: anaesthesia. *Br J Anaesth.* 2011;106(5):617–31.
288. Joffe AM, Aziz MF, Posner KL, Duggan LV, Mincer SL, Domino KB. Management of difficult tracheal intubation: a closed claims analysis. *Anesthesiology.* 2019;131(4):818–29.
289. Greenland KB, Acott C, Segal R, Goulding G, Riley RH, Merry AF. Emergency surgical airway in life-threatening acute airway emergencies-why are we so reluctant to do it? *Anaesth Intensive Care.* 2011;39(4):578–84.

290. Berwick RJ, Gauntlett W, Silverio SA, Wallace H, Mercer S, Brown JM, et al. A mixed-methods pilot study to evaluate a collaborative anaesthetic and surgical training package for emergency surgical cricothyroidotomy. *Anaesth Intensive Care*. 2019;47(4):357–67.
291. Kristensen MS, Teoh WH, Baker PA. Percutaneous emergency airway access; prevention, preparation, technique and training. *Br J Anaesth*. 2015;114(3):357–61.
292. Zasso FB, Perelman VS, Ye XY, Melvin M, Wild E, Tavares W, et al. Effects of prior exposure to a visual airway cognitive aid on decision-making in a simulated airway emergency: a randomised controlled study. *Eur J Anaesthesiol*. 2021;38(8):831–8.
293. Law JA, Duggan LV, Asselin M, Baker P, Crosby E, Downey A, et al. Canadian Airway Focus Group updated consensus-based recommendations for management of the difficult airway: part 2. Planning and implementing safe management of the patient with an anticipated difficult airway. *Can J Anaesth*. 2021;68(9):1405–36.
294. Potnuru P, Artine CA, Hagberg CA. The lost airway. *Anesthesiol Clin*. 2020;38(4):875–88.
295. Dixit A, Ramaswamy KK, Perera S, Sukumar V, Frerk C. Impact of change in head and neck position on ultrasound localisation of the cricothyroid membrane: an observational study. *Anesthesia*. 2019;74(1):29–32.
296. Zasso FB, You-Ten KE, Ryu M, Losyeva K, Tanwani J, Siddiqui N. Complications of cricothyroidotomy versus tracheostomy in emergency surgical airway management: a systematic review. *BMC Anesthesiol*. 2020;20(1):216.
297. DeVore EK, Redmann A, Howell R, Khosla S. Best practices for emergency surgical airway: A systematic review. *Laryngoscope Investig Otolaryngol*. 2019;4(6):602–8.
298. McNiven ND, Pracy JP, McGrath BA, Robson AK. The role of Scalpel-bougie cricothyroidotomy in managing emergency Front of Neck Airway access. A review and technical update for ENT surgeons. *Clin Otolaryngol*. 2018;43(3):791–4.
299. Langvad S, Hyldmo PK, Nakstad AR, Vist GE, Sandberg M. Emergency cricothyrotomy-a systematic review. *Scand J Trauma Resusc Emerg Med*. 2013;21:43.
300. Timmermann A, Chrimis N, Hagberg CA. Need to consider human factors when determining first-line technique for emergency front-of-neck access. *Br J Anaesth*. 2016;117(1):5–7.
301. Onrubia X, Frova G, Sorbello M. Front of neck access to the airway: a narrative review. *Trends in Anesthesia and Critical Care*. 2018;22:45–55.
302. Greenland KB, Bradley WPL, Chapman GA, Goulding G, Irwin MG. Emergency front-of-neck access: scalpel or cannula-and the parable of Buridan's ass†. *Br J Anaesth*. 2017;118(6):811–4.
303. Morton S, Avery P, Kua J, O'Meara M. Success rate of pre-hospital emergency front-of-neck access (FONA): a systematic review and meta-analysis. *Br J Anaesth*. 2023;130(5):636–44.
304. Hubble MW, Wilfong DA, Brown LH, Hertelendy A, Benner RW. A meta-analysis of prehospital airway control techniques part II: alternative airway devices and cricothyrotomy success rates. *Prehosp Emerg Care*. 2010;14(4):515–30.
305. Niven AS, Doerschug KC. Techniques for the difficult airway. *Curr Opin Crit Care*. 2013;19(1):9–15.
306. Lockey D, Crewdson K, Weaver A, Davies G. Observational study of the success rates of intubation and failed intubation airway rescue techniques in 7256 attempted intubations of trauma patients by pre-hospital physicians. *Br J Anaesth*. 2014;113(2):220–5.
307. Law JA. Deficiencies in locating the cricothyroid membrane by palpation: We can't and the surgeons can't, so what now for the emergency surgical airway? *Can J Anaesth*. 2016;63(7):791–6.
308. Schaumann N, Lorenz V, Schellongowski P, Staudinger T, Locker GJ, Burgmann H, et al. Evaluation of Seldinger technique emergency cricothyroidotomy versus standard surgical cricothyroidotomy in 200 cadavers. *Anesthesiology*. 2005;102(1):7–11.
309. Pracy JP, Brennan L, Cook TM, Hartle AJ, Marks RJ, McGrath BA, et al. Surgical intervention during a Can't intubate Can't Oxygenate (CICO) event: emergency front-of-neck airway (FONA)? *Br J Anaesth*. 2016;117(4):426–8.
310. Cook TM, Woodall N, Frerk C. A national survey of the impact of NAP4 on airway management practice in United Kingdom hospitals: closing the safety gap in anaesthesia, intensive care and the emergency department. *Br J Anaesth*. 2016;117(2):182–90.
311. Booth AWG, Vidhani K. Human factors can't intubate can't oxygenate (CICO) bundle is more important than needle versus scalpel debate. *Br J Anaesth*. 2017;118(3):466–8.
312. Chang JE, Kim H, Won D, Lee JM, Kim TK, Min SW, et al. Comparison of the conventional downward and modified upward laryngeal handshake techniques to identify the cricothyroid membrane: a randomized, comparative study. *Anesth Analg*. 2021;133(5):1288–95.
313. Drew T, McCaul CL. Laryngeal handshake technique in locating the cricothyroid membrane: a non-randomised comparative study. *Br J Anaesth*. 2018;121(5):1173–8.
314. Fennelly P, Drew T, Husarova V, Duggan M, McCaul CL. Emergency cricothyroidotomy: an observational study to estimate optimal incision position and length. *Br J Anaesth*. 2019;122(2):263–8.
315. Hung KC, Chen IW, Lin CM, Sun CK. Comparison between ultrasound-guided and digital palpation techniques for identification of the cricothyroid membrane: a meta-analysis. *Br J Anaesth*. 2021;126(1):e9–11.
316. Austin DR, Chang MG, Bittner EA. Use of handheld point-of-care ultrasound in emergency airway management. *Chest*. 2021;159(3):1155–65.
317. Bowness J, Teoh WH, Kristensen MS, Dalton A, Saint-Grant AL, Taylor A, et al. A marking of the cricothyroid membrane with extended neck returns to correct position after neck manipulation and repositioning. *Acta Anaesthesiol Scand*. 2020;64(10):1422–5.
318. Rai Y, You-Ten E, Zasso F, De Castro C, Ye XY, Siddiqui N. The role of ultrasound in front-of-neck access for cricothyroid membrane identification: a systematic review. *J Crit Care*. 2020;60:161–8.
319. Kristensen MS, Teoh WH. Ultrasound identification of the cricothyroid membrane: the new standard in preparing for front-of-neck airway access. *Br J Anaesth*. 2021;126(1):22–7.
320. Choi J, Anderson TN, Sheira D, Sousa J, Borghi JA, Spain DA, et al. The need to routinely convert emergency cricothyroidotomy to tracheostomy: a systematic review and meta-analysis. *J Am Coll Surg*. 2022;234(5):947–52.
321. Talving P, DuBose J, Inaba K, Demetriades D. Conversion of emergent cricothyrotomy to tracheotomy in trauma patients. *Arch Surg*. 2010;145(1):87–91.
322. Myatra SN, Kalkundre RS, Divatia JV. Optimizing education in difficult airway management: meeting the challenge. *Curr Opin Anaesthesiol*. 2017;30(6):748–54.
323. You-Ten KE, Wong C, Arzola C, Cheung J, Friedman Z, Perelman S, et al. Role of contextualizing a crisis scenario on the performance of a cricothyrotomy procedural task. *Can J Anaesth*. 2015;62(10):1104–13.
324. Lemay F, Asselin M, Labrecque P. Leadership and teaching in airway management. *Can J Anaesth*. 2021;68(9):1317–23.
325. Carney N, Totten AM, Cheney T, Jungbauer R, Neth MR, Weeks C, et al. Prehospital airway management: a systematic review. *Prehosp Emerg Care*. 2022;26(5):716–27.
326. Denninghoff KR, Nuño T, Pauls Q, Yeatts SD, Silbergbeit R, Palesch YY, et al. Prehospital intubation is associated with

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

- favorable outcomes and lower mortality in ProTECT III. *Prehosp Emerg Care.* 2017;21(5):539–44.
327. Gaither JB, Spaite DW, Stoltz U, Ennis J, Mosier J, Sakles JJ. Prevalence of difficult airway predictors in cases of failed prehospital endotracheal intubation. *J Emerg Med.* 2014;47(3):294–300.
328. Carlson JN, Colella MR, Daya MR, J De Maio V, Nawrocki P, Nikolla DA, et al. Prehospital cardiac arrest airway management: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):54–63.
329. Braithwaite S, Stephens C, Remick K, Barrett W, Guyette FX, Levy M, et al. Prehospital trauma airway management: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):64–71.
330. Lockey DJ, Crewdson K, Davies G, Jenkins B, Klein J, Laird C, et al. AAGBI: safer pre-hospital anaesthesia 2017: association of anaesthetists of Great Britain and Ireland. *Anaesthesia.* 2017;72(3):379–90.
331. Morton S, Dawson J, Wareham G, Broomhead R, Sherren P. The prehospital emergency anaesthetic in 2022. *Air Med J.* 2022;41(6):530–5.
332. Freund Y, Duchateau FX, Devaud ML, Ricard-Hibon A, Juvin P, Mantz J. Factors associated with difficult intubation in prehospital emergency medicine. *Eur J Emerg Med.* 2012;19(5):304–8.
333. Lockey DJ, Avery P, Harris T, Davies GE, Lossius HM. A prospective study of physician pre-hospital anaesthesia in trauma patients: oesophageal intubation, gross airway contamination and the 'quick look' airway assessment. *BMC Anesthesiol.* 2013;13(1):21.
334. Breckwoldt J, Klemstein S, Brunne B, Schnitzer L, Mochmann HC, Arntz HR. Difficult prehospital endotracheal intubation - predisposing factors in a physician based EMS. *Resuscitation.* 2011;82(12):1519–24.
335. Carlson JN, Hostler D, Guyette FX, Pinchak M, Martin-Gill C. Derivation and Validation of The Prehospital Difficult Airway IdentificationTool (PreDAIT): a predictive model for difficult intubation. *West J Emerg Med.* 2017;18(4):662–72.
336. Jarvis JL, Lyng JW, Miller BL, Perlmutter MC, Abraham H, Sahni R. Prehospital drug assisted airway management: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):42–53.
337. Dorsett M, Panchal AR, Stephens C, Farca A, Leggio W, Galton C, et al. Prehospital airway management training and education: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):3–13.
338. Binks MJ, Holyoak RS, Melhuish TM, Vlok R, Bond E, White LD. Apneic oxygenation during intubation in the emergency department and during retrieval: a systematic review and meta-analysis. *Am J Emerg Med.* 2017;35(10):1542–6.
339. Pavlov I, Medrano S, Weingart S. Apneic oxygenation reduces the incidence of hypoxemia during emergency intubation: a systematic review and meta-analysis. *Am J Emerg Med.* 2017;35(8):1184–9.
340. Spaite DW, Hu C, Bobrow BJ, Chikani V, Barnhart B, Gaither JB, et al. The effect of combined out-of-hospital hypotension and hypoxia on mortality in major traumatic brain injury. *Ann Emerg Med.* 2017;69(1):62–72.
341. Crewdson K, Lockey D, Voelckel W, Temesvari P, Lossius HM, Group EMW. Best practice advice on pre-hospital emergency anaesthesia & advanced airway management. *Scand J Trauma Resusc Emerg Med.* 2019;27(1):6.
342. Benoit JL, Prince DK, Wang HE. Mechanisms linking advanced airway management and cardiac arrest outcomes. *Resuscitation.* 2015;93:124–7.
343. Jeong S, Ahn KO, Shin SD. The role of prehospital advanced airway management on outcomes for out-of-hospital car- diac arrest patients: a meta-analysis. *Am J Emerg Med.* 2016;34(11):2101–6.
344. Benoit JL, Gerecht RB, Steuerwald MT, McMullan JT. Endotracheal intubation versus supraglottic airway placement in out-of-hospital cardiac arrest: a meta-analysis. *Resuscitation.* 2015;93:20–6.
345. Fouche PF, Simpson PM, Bendall J, Thomas RE, Cone DC, Doi SA. Airways in out-of-hospital cardiac arrest: systematic review and meta-analysis. *Prehosp Emerg Care.* 2014;18(2):244–56.
346. Bossers SM, Schwarte LA, Loer SA, Twisk JW, Boer C, Schober P. Experience in prehospital endotracheal intubation significantly influences mortality of patients with severe traumatic brain injury: a systematic review and meta-analysis. *PLoS One.* 2015;10(10):e0141034.
347. Anderson J, Ebeid A, Stallwood-Hall C. Pre-hospital tracheal intubation in severe traumatic brain injury: a systematic review and meta-analysis. *Br J Anaesth.* 2022;129(6):977–84.
348. Kovacs G, Sowers N. Airway management in trauma. *Emerg Med Clin North Am.* 2018;36(1):61–84.
349. Counts CR, Benoit JL, McClelland G, DuCanto J, Weekes L, Latimer A, et al. Novel technologies and techniques for prehospital airway management: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):129–36.
350. Suppan L, Tramèr MR, Niquille M, Grosgeurin O, Marti C. Alternative intubation techniques vs Macintosh laryngoscopy in patients with cervical spine immobilization: systematic review and meta-analysis of randomized controlled trials. *Br J Anaesth.* 2016;116(1):27–36.
351. Pourmand A, Terrebonne E, Gerber S, Shipley J, Tran QK. Efficacy of video laryngoscopy versus direct laryngoscopy in the prehospital setting: a systematic review and meta-analysis. *Prehosp Disaster Med.* 2022;1:1–11.
352. Hayes-Bradley C, Gemal H, Miller M, Ware S. Describing the challenges of prehospital rapid sequence intubation by macintosh blade video laryngoscopy recordings. *Prehosp Disaster Med.* 2022;37(4):485–91.
353. Burgess MB, Schauer SG, Hood RL, De Lorenzo RA. The difficult airway redefined. *Prehosp Disaster Med.* 2022;37(6):723–6.
354. Sultan P, Carvalho B, Rose BO, Cregg R. Endotracheal tube cuff pressure monitoring: a review of the evidence. *J Perioper Pract.* 2011;21(11):379–86.
355. Pluijms WA, van Mook WN, Wittekamp BH, Bergmans DC. Postextubation laryngeal edema and stridor resulting in respiratory failure in critically ill adult patients: updated review. *Crit Care.* 2015;19(1):295.
356. Thiruvenkata Rajan V, Van Wijk RM, Rajbhoj A. Cranial nerve injuries with supraglottic airway devices: a systematic review of published case reports and series. *Anaesthesia.* 2015;70(3):344–59.
357. Kang JE, Oh CS, Choi JW, Son IS, Kim SH. Postoperative pharyngolaryngeal adverse events with laryngeal mask airway (LMA Supreme) in laparoscopic surgical procedures with cuff pressure limiting 25 cmH2O: prospective, blind, and randomised study. *ScientificWorldJournal.* 2014;2014:709801.
358. Haas CF, Eakin RM, Konkle MA, Blank R. Endotracheal tubes: old and new. *Respir Care.* 2014;59(6):933–52.
359. Grant T. Do current methods for endotracheal tube cuff inflation create pressures above the recommended range? A review of the evidence. *J Perioper Pract.* 2013;23(12):292–5.
360. Hockey CA, van Zundert AA, Paratz JD. Does objective measurement of tracheal tube cuff pressures minimise adverse effects and maintain accurate cuff pressures? A systematic review and meta-analysis. *Anaesth Intensive Care.* 2016;44(5):560–70.
361. Schalk R, Seeger FH, Mutlak H, Schweigkofler U, Zacharowski K, Peter N, et al. Complications associated with the prehospital use of laryngeal tubes-a systematic analysis of

- risk factors and strategies for prevention. *Resuscitation*. 2014;85(11):1629–32.
362. Bick E, Bailes I, Patel A, Brain AI. Fewer sore throats and a better seal: why routine manometry for laryngeal mask airways must become the standard of care. *Anaesthesia*. 2014;69(12):1304–8.
363. Peters JH, Hoogerwerf N. Prehospital endotracheal intubation; need for routine cuff pressure measurement? *Emerg Med J*. 2013;30(10):851–3.
364. Hensel M, Güldenpfennig T, Schmidt A, Krumm M. [Continuous cuff pressure measurement during laryngeal mask anesthesia : An obligatory measure to avoid postoperative complications]. *Anaesthesia*. 2016;65(5):346–52.
365. Maertens B, Lin F, Chen Y, Rello J, Lathyris D, Blot S. Effectiveness of continuous cuff pressure control in preventing ventilator-associated pneumonia: a systematic review and meta-analysis of randomized controlled trials. *Crit Care Med*. 2022;50(10):1430–9.
366. Nseir S, Lorente L, Ferrer M, Rouzé A, Gonzalez O, Bassi GL, et al. Continuous control of tracheal cuff pressure for VAP prevention: a collaborative meta-analysis of individual participant data. *Ann Intensive Care*. 2015;5(1):43.
367. Rouzé A, Nseir S. Continuous control of tracheal cuff pressure for the prevention of ventilator-associated pneumonia in critically ill patients: where is the evidence? *Curr Opin Crit Care*. 2013;19(5):440–7.
368. Rackley CR. Monitoring during mechanical ventilation. *Respir Care*. 2020;65(6):832–46.
369. Hess DR, Altobelli NP. Tracheostomy tubes. *Respir Care*. 2014;59(6):956–71, discussion 971-973.
370. Carhart E, Stuck LH, Salzman JG. Achieving a safe endotracheal tube cuff pressure in the prehospital setting: is it time to revise the standard cuff inflation practice? *Prehosp Emerg Care*. 2016;20(2):273–7.
371. Krieger M, Alflein C, Eisel J, Ott T, Piepho T, Noppens RR. Evaluation of the optimal cuff volume and cuff pressure of the revised laryngeal tube LTS-Dün surgical patients. *BMC Anesthesiol*. 2017;17(1):19.
372. El-Orbany M, Salem MR. Endotracheal tube cuff leaks: causes, consequences, and management. *Anesth Analg*. 2013;117(2):428–34.
373. Beydon L, Gourges M, Talec P. [Endotracheal tube cuff and nitrous oxide: bench evaluation and assessment of clinical practice]. *Ann Fr Anesth Reanim*. 2011;30(9):679–84.
374. Maggiore SM, Battilana M, Serano L, Petrini F. Ventilatory support after extubation in critically ill patients. *Lancet Respir Med*. 2018;6(12):948–62.
375. Gómez-Ríos MÁ, Abad-Gurumeta A, Casans-Francés R, Esquinas AM. Safe extubation procedure of the difficult airway: think twice, act wise. *Minerva Anestesiol*. 2020;86(8):802–4.
376. Parotto M, Cooper RM, Behringer EC. Extubation of the Challenging or Difficult Airway. *Curr Anesthesiol Rep*. 2020;1–7.
377. Juang J, Cordoba M, Ciaramella A, Xiao M, Goldfarb J, Bayter JE, et al. Incidence of airway complications associated with deep extubation in adults. *BMC Anesthesiol*. 2020;20(1):274.
378. Benham-Hermetz J, Mitchell V. Safe tracheal extubation after general anaesthesia. *BJA Education*. 2021;21(12):446–54.
379. Cavallone LF, Vannucci A. Review article: Extubation of the difficult airway and extubation failure. *Anesth Analg*. 2013;116(2):368–83.
380. Mitchell V, Cooper R. Extubation. In: Kristensen MS, Cook T, editors. *Core Topics in Airway Management*. 3 ed. Cambridge: Cambridge University Press; 2020. p. 177–84.
381. Kellner DB, Urman RD, Greenberg P, Brovman EY. Analysis of adverse outcomes in the post-anesthesia care unit based on anesthesia liability data. *J Clin Anesth*. 2018;50:48–56.
382. Torrini F, Gendreau S, Morel J, Carteaux G, Thille AW, Antonelli M, et al. Prediction of extubation outcome in critically ill patients: a systematic review and meta-analysis. *Crit Care*. 2021;25(1):391.
383. Sturgess DJ, Greenland KB, Senthuran S, Ajvadi FA, van Zundert A, Irwin MG. Tracheal extubation of the adult intensive care patient with a predicted difficult airway - a narrative review. *Anaesthesia*. 2017;72(2):248–61.
384. Cook TM, Woodall N, Frerk C, Project FNA. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 1: anaesthesia. *Br J Anaesth*. 2011;106(5):617–31.
385. Bösel J. Who is safe to extubate in the neuroscience intensive care unit? *Semin Respir Crit Care Med*. 2017;38(6):830–9.
386. Parotto M., Ellard L. Extubation following anesthesia. UpToDate. Retrieved November 2021.
387. Nwakanma CC, Wright BJ. Extubation in the emergency department and resuscitative unit setting. *Emerg Med Clin North Am*. 2019;37(3):557–68.
388. Kuriyama A, Jackson JL, Kamei J. Performance of the cuff leak test in adults in predicting post-extubation airway complications: a systematic review and meta-analysis. *Crit Care*. 2020;24(1):640.
389. Hyzy RC. Extubation management in the adult intensive care unit. UpToDate. Retrieved November 2021.
390. Sutherasan Y, Theerawit P, Hongphanut T, Kiatboonsri C, Kiatboonsri S. Predicting laryngeal edema in intubated patients by portable intensive care unit ultrasound. *J Crit Care*. 2013;28(5):675–80.
391. Mikaeili H, Yazdchi M, Tarzamni MK, Ansarin K, Ghasemzadeh M. Laryngeal ultrasonography versus cuff leak test in predicting postextubation stridor. *J Cardiovasc Thorac Res*. 2014;6(1):25–8.
392. Ochoa ME, Marín Me C, Frutos-Vivar F, Gordo F, Latour-Pérez J, Calvo E, et al. Cuff-leak test for the diagnosis of upper airway obstruction in adults: a systematic review and meta-analysis. *Intensive Care Med*. 2009;35(7):1171–9.
393. Carvalho H, Verdonck M, Cools W, Geerts L, Forget P, Poelaert J. Forty years of neuromuscular monitoring and postoperative residual curarisation: a meta-analysis and evaluation of confidence in network meta-analysis. *Br J Anaesth*. 2020;125(4):466–82.
394. Carron M, Zarantonello F, Tellaroli P, Ori C. Efficacy and safety of sugammadex compared to neostigmine for reversal of neuromuscular blockade: a meta-analysis of randomized controlled trials. *J Clin Anesth*. 2016;35:1–12.
395. Cammu G. Residual Neuromuscular Blockade and Postoperative Pulmonary Complications: What Does the Recent Evidence Demonstrate? *Curr Anesthesiol Rep*. 2020;1–6.
396. Raval AD, Uyei J, Karabis A, Bash LD, Brull SJ. Incidence of residual neuromuscular blockade and use of neuromuscular blocking agents with or without antagonists: A systematic review and meta-analysis of randomized controlled trials. *J Clin Anesth*. 2020;64:109818.
397. Kuriyama A, Umakoshi N, Sun R. Prophylactic corticosteroids for prevention of postextubation stridor and reintubation in adults: a systematic review and meta-analysis. *Chest*. 2017;151(5):1002–10.
398. Ahn C, Na MK, Choi KS, Lim TH, Jang BH, Kim W, et al. Comparison between multiple doses and single-dose steroids in preventing the incidence of reintubation after extubation among critically ill patients: a network meta-analysis. *J Clin Med*. 2021;10(13).
399. Jaber S, Jung B, Chanques G, Bonnet F, Marret E. Effects of steroids on reintubation and post-extubation stridor in adults: meta-analysis of randomised controlled trials. *Crit Care*. 2009;13(2):R49.
400. Sakae TM, Souza RLP, Brand Ú JCM. Impact of topical airway anesthesia on immediate postoperative cough/bucking:

M.Á. Gómez-Ríos, J.A. Sastre, X. Onrubia-Fuertes et al.

- a systematic review and meta-analysis. *Braz J Anesthesiol.* 2023;73(1):91–100.
401. Peng F, Wang M, Yang H, Yang X, Long M. Efficacy of intracuff lidocaine in reducing coughing on tube: a systematic review and meta-analysis. *J Int Med Res.* 2020;48(2):300060520901872.
402. Yang SS, Wang NN, Postonogova T, Yang GJ, McGillion M, Beique F, et al. Intravenous lidocaine to prevent postoperative airway complications in adults: a systematic review and meta-analysis. *Br J Anaesth.* 2020;124(3):314–23.
403. Clivio S, Putzu A, Tramèr MR. Intravenous lidocaine for the prevention of cough: systematic review and meta-analysis of randomized controlled trials. *Anesth Analg.* 2019;129(5):1249–55.
404. Zhang J, Yu Y, Miao S, Liu L, Gan S, Kang X, et al. Effects of peri-operative intravenous administration of dexmedetomidine on emergence agitation after general anesthesia in adults: a meta-analysis of randomized controlled trials. *Drug Des Devel Ther.* 2019;13:2853–64.
405. Kim JH, Ham SY, Kim DH, Chang CH, Lee JS. Efficacy of single-dose dexmedetomidine combined with low-dose remifentanil infusion for cough suppression compared to high-dose remifentanil infusion: a randomized, controlled, non-inferiority trial. *Int J Med Sci.* 2019;16(3):376–83.
406. Wong TH, Weber G, Abramowicz AE. Smooth extubation and smooth emergence techniques: a narrative review. *Anesthesiol Res Pract.* 2021;2021:8883257.
407. Kalra N, Gupta A, Sood R, Kaur M. Comparison of proseal laryngeal mask airway with the I-Gel supraglottic airway during the bailey manoeuvre in adult patients undergoing elective surgery. *Turk J Anaesthesiol Reanim.* 2021;49(2):107–13.
408. Tung A, Fergusson NA, Ng N, Hu V, Dormuth C, Griesdale DEG. Medications to reduce emergence coughing after general anaesthesia with tracheal intubation: a systematic review and network meta-analysis. *Br J Anaesth.* 2020.
409. Salim B, Rashid S, Ali MA, Raza A, Khan FA. Effect of Pharmacological Agents Administered for Attenuating the Extubation Response on the Quality of Extubation: A Systematic Review. *Cureus.* 2019;11(12):e6427.
410. Artime CA, Hagberg CA. Tracheal extubation. *Respir Care.* 2014;59(6):991–1002.
411. Duggan LV, Law JA, Murphy MF. Brief review: Supplementing oxygen through an airway exchange catheter: efficacy, complications, and recommendations. *Can J Anaesth.* 2011;58(6):560–8.
412. Mort TC. Continuous airway access for the difficult extubation: the efficacy of the airway exchange catheter. *Anesth Analg.* 2007;105(5):1357–62.
413. Furyk C, Walsh ML, Kaliaperumal I, Bentley S, Hattingh C. Assessment of the reliability of intubation and ease of use of the Cook Staged Extubation Set—an observational study. *Anaesth Intensive Care.* 2017;45(6):695–9.
414. McManus S, Jones L, Anstey C, Senthuran S. An assessment of the tolerability of the Cook staged extubation wire in patients with known or suspected difficult airways extubated in intensive care. *Anaesthesia.* 2018;73(5):587–93.
415. Lu C, Li J, Zhao S, Zhang Y. Efficacy and safety of Cook staged Extubation Set in patients with difficult airway: a systematic review and meta-analysis. *BMC Anesthesiol.* 2023;23(1):232.
416. Corso RM, Sorbello M, Mecugni D, Seligardi M, Piraccini E, Agnolletti V, et al. Safety and efficacy of Staged Extubation Set in patients with difficult airway: a prospective multicenter study. *Minerva Anestesiol.* 2020;86(8):827–34.
417. Mort TC, Braffett BH. Conventional versus video laryngoscopy for tracheal tube exchange: glottic visualization, success rates, complications, and rescue alternatives in the high-risk difficult airway patient. *Anesth Analg.* 2015;121(2):440–8.
418. Yasuda H, Okano H, Mayumi T, Narita C, Onodera Y, Nakane M, et al. Post-extubation oxygenation strategies in acute respiratory failure: a systematic review and network meta-analysis. *Crit Care.* 2021;25(1):135.
419. Zhou X, Yao S, Dong P, Chen B, Xu Z, Wang H. Preventive use of respiratory support after scheduled extubation in critically ill medical patients—a network meta-analysis of randomized controlled trials. *Crit Care.* 2020;24(1):370.
420. Bajaj A, Rathor P, Sehgal V, Shetty A. Efficacy of noninvasive ventilation after planned extubation: A systematic review and meta-analysis of randomized controlled trials. *Heart Lung.* 2015;44(2):150–7.
421. Glossop AJ, Shephard N, Bryden DC, Mills GH. Non-invasive ventilation for weaning, avoiding reintubation after extubation and in the postoperative period: a meta-analysis. *Br J Anaesth.* 2012;109(3):305–14.
422. Lundstrøm LH, Møller AM, Rosenstock C, Astrup G, Gätke MR, Wetterslev J, et al. A documented previous difficult tracheal intubation as a prognostic test for a subsequent difficult tracheal intubation in adults. *Anaesthesia.* 2009;64(10):1081–8.
423. Schechtman SA, Flori HR, Thatcher AL, Almendras G, Robell SE, Healy DW, et al. The difficult airway navigator: development and implementation of a health care system's approach to difficult airway documentation utilizing the electronic health record. *A A Pract.* 2021;15(5):e01455.
424. Feinleib J, Foley L, Mark L. What we all should know about our patient's airway: difficult airway communications, database registries, and reporting systems registries. *Anesthesiol Clin.* 2015;33(2):397–413.
425. Zaouter C, Calderon J, Hemmerling TM. Videolaryngoscopy as a new standard of care. *Br J Anaesth.* 2015;114(2):181–3.
426. Sajayan A, Nair A, McNarry AF, Mir F, Ahmad I, El-Boghdadly K. Analysis of a national difficult airway database. *Anaesthesia.* 2022;77(10):1081–8.
427. Matava C, Caldeira-Kulkarni M, Chisholm J. Improved difficult airway documentation using structured notes in Anesthesia Information Management Systems. *Can J Anaesth.* 2020;67(5):625–7.
428. McGrath BA, Calder N, Laha S, Perks A, Chaudry I, Bates L, et al. Reduction in harm from tracheostomy-related patient safety incidents following introduction of the National Tracheostomy Safety Project: our experience from two hundred and eighty-seven incidents. *Clin Otolaryngol.* 2013;38(6):541–5.
429. McNarry AF, M Cook T, Baker PA, O'Sullivan EP. The Airway Lead: opportunities to improve institutional and personal preparedness for airway management. *Br J Anaesth.* 2020;125(1):e22–4.
430. Smith C, McNarry AF. Airway leads and airway response teams: improving delivery of safer airway management? *Curr Anesthesiol Rep.* 2020;1:8.
431. Gurses AP, Marsteller JA, Ozok AA, Xiao Y, Owens S, Pronovost PJ. Using an interdisciplinary approach to identify factors that affect clinicians' compliance with evidence-based guidelines. *Crit Care Med.* 2010;38 8 Suppl:S282–91.
432. Paksaite P, Crosskey J, Sula E, West C, Watson M. A systematic review using the Theoretical Domains Framework to identify barriers and facilitators to the adoption of prescribing guidelines. *Int J Pharm Pract.* 2021;29(1):3–11.
433. Gurses AP, Murphy DJ, Martinez EA, Berenholtz SM, Pronovost PJ. A practical tool to identify and eliminate barriers to compliance with evidence-based guidelines. *Comm J Qual Patient Saf.* 2009;35(10):526–32, 485.
434. Jordan P, Mpasa F, Ten Ham-Baloyi W, Bowers C. Implementation strategies for guidelines at ICUs: a systematic review. *Int J Health Care Qual Assur.* 2017;30(4):358–72.
435. Mitchell RJ, Williamson AM, Molesworth B, Chung AZ. A review of the use of human factors classification frameworks that

- identify causal factors for adverse events in the hospital setting. *Ergonomics*. 2014;57(10):1443–72.
436. Weigel WA. Redesigning an airway cart using lean methodology. *J Clin Anesth*. 2016;33:273–82.
437. Bjurström MF, Persson K, Sturesson LW. Availability and organization of difficult airway equipment in Swedish hospitals: a national survey of anaesthesiologists. *Acta Anaesthesiol Scand*. 2019;63(10):1313–20.
438. Greenland KB. Introducing airway leads in Australia. Australian and New Zealand College of Anaesthetists. 2019;22–3.
439. Baker PA, Behringer EC, Feinleib J, Foley LJ, Mosier J, Roth P, Wali A, O'Sullivan EP. Formation of an Airway Lead Network: an essential patient safety initiative. *Br J Anaesth*. 2022;128(2):225–9.
440. Mark L, Lester L, Cover R, Herzer K. A decade of difficult airway response team: lessons learned from a hospital-wide difficult airway response team program. *Crit Care Clin*. 2018;34(2):239–51.
441. Damrose JF, Eropkin W, Ng S, Cale S, Banerjee S. The critical response team in airway emergencies. *Perm J*. 2019;23.
442. Tankard KA, Sharifpour M, Chang MG, Bittner EA. Design and implementation of airway response teams to improve the practice of emergency airway management. *J Clin Med*. 2022;11(21).
443. Pandit JJ, Popat MT, Cook TM, Wilkes AR, Groom P, Cooke H, et al. The Difficult Airway Society' ADEPT' guidance on selecting airway devices: the basis of a strategy for equipment evaluation. *Anesthesia*. 2011;66(8):726–37.
444. Athanassoglou V, O'Sullivan EP, van Zundert A, Pandit JJ. New guidelines for research in airway device evaluation: time for an updated approach (ADEPT-2) to the Difficult Airway Society's' ADEPT' strategy? *J Clin Monit Comput*. 2023;37(2):345–50.
445. Marshall SD, Chrimes N. Time for a breath of fresh air: Rethinking training in airway management. *Anesthesia*. 2016;71(11):1259–64.
446. Baker PA, Feinleib J, O'Sullivan EP. Is it time for airway management education to be mandatory? *Br J Anaesth*. 2016;117 Suppl 1:i13–6.
447. Howard M, Noppens R, Gonzalez N, Jones PM, Payne SM. Seven years on from the Canadian Airway Focus Group Difficult Airway Guidelines: an observational survey. *Can J Anaesth*. 2021;68(9):1331–6.
448. Brown W, Santhosh L, Brady AK, Denson JL, Niroula A, Pugh ME, et al. A call for collaboration and consensus on training for endotracheal intubation in the medical intensive care unit. *Crit Care*. 2020;24(1):621.
449. Armstrong L, Harding F, Critchley J, McNarry AF, Myatra SN, Cooper R, et al., Group WAMME. An international survey of airway management education in 61 countries. *Br J Anaesth*. 2020;125(1):e54–60.
450. Lindkaer Jensen NH, Cook TM, Kelly FE. A national survey of practical airway training in UK anaesthetic departments. Time for a national policy? *Anesthesia*. 2016;71(11):1273–9.
451. Kelly FE, Frerk C, Bailey CR, Cook TM, Ferguson K, Flin R, et al. Human factors in anaesthesia: a narrative review. *Anesthesia*. 2023;78(4):479–90.
452. Brydges R, Hatala R, Zendejas B, Erwin PJ, Cook DA. Linking simulation-based educational assessments and patient-related outcomes: a systematic review and meta-analysis. *Acad Med*. 2015;90(2):246–56.
453. Hatala R, Cook DA, Zendejas B, Hamstra SJ, Brydges R. Feedback for simulation-based procedural skills training: a meta-analysis and critical narrative synthesis. *Adv Health Sci Educ Theory Pract*. 2014;19(2):251–72.
454. Cook DA, Brydges R, Zendejas B, Hamstra SJ, Hatala R. Mastery learning for health professionals using technology-enhanced simulation: a systematic review and meta-analysis. *Acad Med*. 2013;88(8):1178–86.
455. Grande B, Kolbe M, Biro P. Difficult airway management and training: simulation, communication, and feedback. *Curr Opin Anaesthesiol*. 2017;30(6):743–7.
456. Jensen JK, Wisborg T. Training and assessment of anaesthesiologist skills: The contrasting groups method and mastery learning levels. *Acta Anaesthesiol Scand*. 2018;62(6):742–3.
457. Donoghue A, Navarro K, Diederich E, Auerbach M, Cheng A. Deliberate practice and mastery learning in resuscitation education: a scoping review. *Resusc Plus*. 2021;6:100137.
458. Petrosoniak A, Lu M, Gray S, Hicks C, Sherbino J, McGowan M, et al. Perfecting practice: a protocol for assessing simulation-based mastery learning and deliberate practice versus self-guided practice for bougie-assisted cricothyroidotomy performance. *BMC Med Educ*. 2019;19(1):100.
459. Nielsen RP, Nikolajsen L, Paltved C, Aagaard R. Effect of simulation-based team training in airway management: a systematic review. *Anesthesia*. 2021;76(10):1404–15.
460. Buljac-Samardzic M, Doekhie KD, van Wijngaarden JDH. Interventions to improve team effectiveness within health care: a systematic review of the past decade. *Hum Resour Health*. 2020;18(1):2.
461. Merriel A, Ficquet J, Barnard K, Kunutsor SK, Soar J, Lenguerand E, et al. The effects of interactive training of healthcare providers on the management of life-threatening emergencies in hospital. *Cochrane Database Syst Rev*. 2019;9:CD012177.
462. Kuzovlev A, Monsieurs KG, Gilfoyle E, Finn J, Greif R. Resuscitation ElaTTfotILCo. The effect of team and leadership training of advanced life support providers on patient outcomes: a systematic review. *Resuscitation*. 2021;160:126–39.
463. Brazil V. Translational simulation: not 'where?' but 'why?' A functional view of in situ simulation. *Advances in Simulation*. 2017;2(1):20.
464. Munzer BW, Bassin BS, Peterson WJ, Tucker RV, Doan J, Harvey C, et al. In-situ simulation use for rapid implementation and process improvement of COVID-19 airway management. *West J Emerg Med*. 2020;21(6):99–106.
465. Levett-Jones T, Lapkin S. The effectiveness of debriefing in simulation-based learning for health professionals: a systematic review. *JBI Libr Syst Rev*. 2012;10(51):3295–337.
466. Johnson TJ, Millinchamp FJ, Kelly FE. Use of a team immediate debrief tool to improve staff well-being after potentially traumatic events. *Anesthesia*. 2021;76(7):1001–2.
467. Couper K, Salman B, Soar J, Finn J, Perkins GD. Debriefing to improve outcomes from critical illness: a systematic review and meta-analysis. *Intensive Care Med*. 2013;39(9):1513–23.
468. Siu LW, Boet S, Borges BC, Bruppacher HR, LeBlanc V, Naik VN, et al. High-fidelity simulation demonstrates the influence of anesthesiologists' age and years from residency on emergency cricothyroidotomy skills. *Anesth Analg*. 2010;111(4):955–60.
469. Karamchandani K, Wheelwright J, Yang AL, Westphal ND, Khanna AK, Myatra SN. Emergency airway management outside the operating room: current evidence and management strategies. *Anesth Analg*. 2021;133(3):648–62.
470. Petersen JA, Bray L, Østergaard D. Continuing professional development for anesthesiologists: a systematic review protocol. *Acta Anaesthesiol Scand*. 2022;66(1):152–5.
471. Clark CA, Mester RA, Redding AT, Wilson DA, Zeiler LL, Jones WR, et al. Emergency subglottic airway training and assessment of skills retention of attending anesthesiologists with simulation mastery-based learning. *Anesth Analg*. 2022;135(1):143–51.
472. Watkins SC, Roberts DA, Boulet JR, McEvoy MD, Weinger MB. Evaluation of a simpler tool to assess nontechnical skills during simulated critical events. *Simul Healthc*. 2017;12(2):69–75.